

**This Page Is Inserted by IFW Operations
and is not a part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- **BLACK BORDERS**
- **TEXT CUT OFF AT TOP, BOTTOM OR SIDES**
- **FADED TEXT**
- **ILLEGIBLE TEXT**
- **SKEWED/SLANTED IMAGES**
- **COLORLED PHOTOS**
- **BLACK OR VERY BLACK AND WHITE DARK PHOTOS**
- **GRAY SCALE DOCUMENTS**


IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/53, 9/02, C12Q 1/68, C12N 1/21, C07K 16/40, A61K 38/44	A1	(11) International Publication Number: WO 98/20137 (43) International Publication Date: 14 May 1998 (14.05.98)
(21) International Application Number: PCT/US97/20681 (22) International Filing Date: 5 November 1997 (05.11.97) (30) Priority Data: 08/745,934 7 November 1996 (07.11.96) US (71) Applicant (for all designated States except US): INCTYE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): HILLMAN, Jennifer, L. [US/US]; 230 Monroe Drive #12, Mountain View, CA 94040 (US). HAWKINS, Phillips, R. [US/US]; 750 N. Shoreline Boulevard #96, Mountain View, CA 94034 (US). (74) Agent: BILLINGS, Lucy, J.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US).		(81) Designated States: AT, AU, BR, CA, CH, CN, DE, DK, ES, FI, GB, IL, JP, KR, MX, NO, NZ, RU, SE, SG, US, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: HUMAN SQUALENE EPOXIDASE 		
(57) Abstract <p>The present invention provides a human squalene epoxidase (HSQEP) and polynucleotides which identify and encode HSQEP. The invention also provides genetically engineered expression vectors and host cells comprising the nucleic acid sequences encoding HSQEP and a method for producing HSQEP. The invention also provides for the use of HSQEP and agonists, antibodies, or antagonists specifically binding HSQEP, in the prevention and treatment of diseases associated with expression of HSQEP. Additionally, the invention provides for the use of antisense molecules to polynucleotides encoding HSQEP for the treatment of diseases associated with the expression of HSQEP. The invention also provides diagnostic assays which utilize the polynucleotide, or fragments or the complement thereof, and antibodies specifically binding HSQEP.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

HUMAN SQUALENE EPOXIDASE

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of a novel squalene epoxidase and to the use of these sequences in the diagnosis, prevention, and treatment of diseases such as hypercholesterolemia and atherosclerosis.

5

BACKGROUND ART

The de novo biosynthesis of cholesterol in the endoplasmic reticulum proceeds by a multistep process. Acetyl coenzyme A (-CoA) is condensed and converted to 3-hydroxy 3-methylglutaryl (HMG)-CoA. HMG-CoA is reduced to mevalonate by the action of HMG-CoA reductase. In consecutive enzyme-catalyzed reactions, mevalonate is converted to isopentenyl pyrophosphate (IPPP), geranyl pyrophosphate (GPP), farnesyl pyrophosphate (FPP) and squalene. Squalene is oxidized to squalene 2,3-epoxide by squalene epoxidase. Rearrangements of squalene 2,3-epoxide lead to lanosterol. Lanosterol is converted to cholesterol in the endoplasmic reticulum in another series of enzymatic reactions.

HMG-CoA reductase is considered one of the major regulatory enzymes in cholesterol biosynthesis. HMG-CoA reductase inhibitors such as lovastatin are widely used to lower plasma cholesterol levels. Since the HMG-CoA reductase-catalyzed production of mevalonate is an early step in the cholesterol biosynthetic pathway, HMG-CoA reductase inhibition depletes many other intermediates of the cholesterol biosynthetic pathway.

Many intermediates in the cholesterol biosynthetic pathway depleted by inhibition of HMG-CoA reductase have additional roles in cellular function. For instance, FPP and FPP-derived geranylgeranyl pyrophosphate (GGPP) covalently modify proteins, heme, and tRNA, and are precursors for biologically important molecules such as dolichols and ubiquinones (Grunler, J. et al. (1994) Biochim. Biophys. Acta 1212:259-277). Posttranslational protein isoprenylation promotes the anchoring of proteins to membranes and serves as a regulatory signal (Glomset, J.A. et al. (1990) Trends Biochem. Sci. 15:139-142). For example, the Rab proteins are a class of small GTP binding proteins which are involved in intracellular vesicle trafficking. Isoprenylated Rab proteins on the surface of vesicles interact with GTPase-activating proteins and specific Rab receptors on the target membrane, leading to membrane

fusion. When cellular isoprenoid synthesis is blocked by lovastatin, Rab proteins that are normally localized in membranes of the endoplasmic reticulum, Golgi apparatus, and endosomes accumulate in the cytosol (Kinsella B.T. et al. (1992) J. Biol. Chem. 267:3940-3945).

5 Defects in protein isoprenylation caused by depletion of FPP and its metabolites may have undesirable biological consequences. The vesicular trafficking of integral membrane proteins is compromised by altered Rab isoprenylation resulting from depletion of intracellular FPP and GGPP. For instance, cystic fibrosis transmembrane conductance regulator (CFTR) function in a primary human airway epithelial cell line is compromised by
10 lovastatin (Shen, B.-Q. et al. (1995) J Biol. Chem. 270:25102-25106). Lovastatin is proposed to disrupt the trafficking of CFTR to the apical plasma membrane by inhibiting the isoprenylation of Rab or Rab-like trafficking proteins. Protein isoprenylation is also important in the maintenance of retinal cytoarchitecture. Lovastatin produces profound dysplastic-like changes in adult rat retinas primarily in the photoreceptor layer (Pittler, S.J. et
15 al. (1995) J. Cell Biol. 130:431-439). This retinal degeneration is traced to defects in protein isoprenylation.

Arteriosclerosis, a generic term for thickening and hardening of the arterial wall, is responsible for the majority of deaths in the United States and most westernized societies. One type of arteriosclerosis is atherosclerosis, the disorder of the larger arteries that underlies
20 most coronary artery disease, aortic aneurysm, and arterial disease of the lower extremities and also plays a major role in cerebrovascular disease. Atherosclerosis is by far the leading cause of death in the United States, both above and below age 65 and in both sexes.

A generally accepted theory for the pathogenesis of atherosclerosis is the reaction to injury hypothesis. According to this idea, the endothelial cells lining the intima are exposed
25 to repeated or continuing insults to their integrity. Injury to the endothelium includes metabolic injury, such as chronic hypercholesterolemia. Reduction of hypercholesterolemia results in a decrease in the progression of atherosclerosis in humans and other primates. Drugs that act primarily by lowering low density lipoprotein (LDL) cholesterol are the current drugs of choice for high-risk patients.

30 It is apparent that, while inhibitors of enzymes at early steps in the cholesterol

biosynthetic pathway are effective cholesterol-lowering therapeutics, such inhibitors also deplete other necessary metabolic intermediates. Inhibitors of enzymes which are further along the cholesterol biosynthetic pathway would therefore provide more desirable cholesterol-lowering therapeutics than the inhibitors of earlier pathway enzymes such as

5 HMG-CoA reductase.

The discovery of polynucleotides encoding human squalene epoxidase, and the molecules themselves, presents opportunities to investigate the regulation and control of the later (post-FPP) cholesterol biosynthetic pathway and to elucidate mechanisms for the reduction of hypercholesterolemia in humans. Discovery of molecules related to squalene

10 epoxidase satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the reduction of LDL cholesterol, a key risk factor in atherosclerosis and coronary heart disease.

DISCLOSURE OF THE INVENTION

The present invention features a human squalene epoxidase hereinafter designated

15 HSQEP and characterized as having chemical and structural homology to squalene epoxidase from rat and mouse. Accordingly, the invention features a substantially purified HSQEP having the amino acid sequence, SEQ ID NO:1.

One aspect of the invention features isolated and substantially purified polynucleotides that encode HSQEP. In a particular aspect, the polynucleotide is the

20 nucleotide sequence of SEQ ID NO:2.

The invention also relates to a polynucleotide sequence comprising the complement of SEQ ID NO:2 or variants thereof. In addition, the invention features polynucleotide sequences which hybridize under stringent conditions to SEQ ID NO:2.

The invention additionally features nucleic acid sequences encoding polypeptides,

25 oligonucleotides, peptide nucleic acids (PNA), fragments, portions or antisense molecules thereof, and expression vectors and host cells comprising polynucleotides that encode HSQEP. The present invention also features antibodies which bind specifically to HSQEP, and pharmaceutical compositions comprising substantially purified HSQEP. The invention also features the use of agonists and antagonists of HSQEP.

30

BRIEF DESCRIPTION OF DRAWINGS

Figures 1A, 1B, 1C, 1D, 1E, 1F and 1G show the amino acid sequence (SEQ ID NO:1) and nucleic acid sequence (SEQ ID NO:2) of HSQEP. The alignment was produced using MacDNASIS PRO™ software (Hitachi Software Engineering Co., Ltd., San Bruno, CA).

Figures 2A, 2B and 2C show the amino acid sequence alignments among HSQEP (SEQ ID NO:1), squalene epoxidase from mouse (GI 1217593; SEQ ID NO:3), and from rat (GI 1083804; SEQ ID NO:4). The alignment was produced using the multisequence alignment program of DNASTAR™ software (DNASTAR Inc, Madison WI).

Figure 3 shows the hydrophobicity plot (MacDNASIS PRO software) for HSQEP, SEQ ID NO: 1; the positive X axis reflects amino acid position, and the negative Y axis, hydrophobicity.

Figure 4 shows the hydrophobicity plot for mouse squalene epoxidase, SEQ ID NO:3.

Figure 5 shows the cholesterol biosynthetic pathway.

MODES FOR CARRYING OUT THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular methodology, protocols, cell lines, vectors, and reagents described as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a", "an", and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, reference to "a host cell" includes a plurality of such host cells, reference to the "antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods, devices,

and materials are now described. All publications mentioned herein are incorporated herein by reference for the purpose of describing and disclosing the cell lines, vectors, and methodologies which are reported in the publications which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not
5 entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

“Nucleic acid sequence” as used herein refers to an oligonucleotide, nucleotide, or polynucleotide, and fragments or portions thereof, and to DNA or RNA of genomic or synthetic origin which may be single- or double-stranded, and represent the sense or antisense
10 strand. Similarly, “amino acid sequence” as used herein refers to an oligopeptide, peptide, polypeptide, or protein sequence and fragments or portions thereof, of a naturally occurring or synthetic molecule.

Where “amino acid sequence” is recited herein to refer to an amino acid sequence of a naturally occurring protein molecule, “amino acid sequence” and like terms, such as
15 “polypeptide” or “protein” are not meant to limit the amino acid sequence to the complete, native amino acid sequence associated with the recited protein molecule.

“Peptide nucleic acid”, as used herein, refers to a molecule which comprises an oligomer to which an amino acid residue, such as lysine, and an amino group have been added. These small molecules, also designated anti-gene agents, stop transcript elongation by
20 binding to their complementary strand of nucleic acid (Nielsen, P.E. et al. (1993) Anticancer Drug Des. 8:53-63).

HSQEP, as used herein, refers to the amino acid sequences of substantially purified HSQEP obtained from any species, particularly mammalian, including bovine, ovine, porcine, equine, and preferably human, from any source whether natural, synthetic,
25 semi-synthetic, or recombinant.

“Consensus”, as used herein, refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, or which has been extended using XL-PCR™ (Perkin Elmer, Norwalk, CT) in the 5' and/or the 3' direction and resequenced, or which has been assembled from the overlapping sequences of more than one Incyte clone using the
30 GELVIEW™ Fragment Assembly system (GCG, Madison, WI), or which has been both

extended and assembled.

A "variant" of HSQEP, as used herein, refers to an amino acid sequence that is altered by one or more amino acids. The variant may have "conservative" changes, wherein a substituted amino acid has similar structural or chemical properties, e.g., replacement of leucine with isoleucine. More rarely, a variant may have "nonconservative" changes, e.g., replacement of a glycine with a tryptophan. Similar minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be substituted, inserted, or deleted without abolishing biological or immunological activity may be found using computer programs well known in the art, for example, DNASTAR software.

A "deletion", as used herein, refers to a change in either amino acid or nucleotide sequence in which one or more amino acid or nucleotide residues, respectively, are absent.

An "insertion" or "addition", as used herein, refers to a change in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid or nucleotide residues, respectively, as compared to the naturally occurring molecule.

A "substitution", as used herein, refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

The term "biologically active", as used herein, refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic HSQEP, or any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The term "agonist", as used herein, refers to a molecule which, when bound to HSQEP, causes a change in HSQEP which modulates the activity of HSQEP. Agonists may include proteins, nucleic acids, carbohydrates, or any other molecules which bind to HSQEP.

The terms "antagonist" or "inhibitor", as used herein, refer to a molecule which, when bound to HSQEP, blocks the biological or immunological activity of HSQEP. Antagonists and inhibitors may include proteins, nucleic acids, carbohydrates, or any other molecules which bind to HSQEP.

The term "modulate", as used herein, refers to a change or an alteration in the

biological activity of HSQEP. Modulation may be an increase or a decrease in protein activity, a change in binding characteristics, or any other change in the biological, functional or immunological properties of HSQEP.

5 The term "mimetic", as used herein, refers to a molecule, the structure of which is developed from knowledge of the structure of HSQEP or portions thereof and, as such, is able to effect some or all of the actions of squalene epoxidase-like molecules.

The term "derivative", as used herein, refers to the chemical modification of a nucleic acid encoding HSQEP or the encoded HSQEP. Illustrative of such modifications would be replacement of hydrogen by an alkyl, acyl, or amino group. A nucleic acid derivative would
10 encode a polypeptide which retains essential biological characteristics of the natural molecule.

The term "substantially purified", as used herein, refers to nucleic or amino acid sequences that are removed from their natural environment, isolated or separated, and are at least 60% free, preferably 75% free, and most preferably 90% free from other components
15 with which they are naturally associated.

"Amplification" as used herein refers to the production of additional copies of a nucleic acid sequence and is generally carried out using polymerase chain reaction (PCR) technologies well known in the art (Dieffenbach, C.W. and G.S. Dveksler (1995) PCR Primer, a Laboratory Manual, Cold Spring Harbor Press, Plainview, NY).

20 The term "hybridization", as used herein, refers to any process by which a strand of nucleic acid binds with a complementary strand through base pairing.

The term "hybridization complex", as used herein, refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen binds between complementary G and C bases and between complementary A and T bases; these hydrogen
25 bonds may be further stabilized by base stacking interactions. The two complementary nucleic acid sequences hydrogen bond in an antiparallel configuration. A hybridization complex may be formed in solution (e.g., C_0t or R_0t analysis) or between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., membranes, filters, chips, pins or glass slides to which cells have been fixed for
30 in situ hybridization).

The terms "complementary" or "complementarity", as used herein, refer to the natural binding of polynucleotides under permissive salt and temperature conditions by base-pairing. For example, for the sequence "A-G-T" binds to the complementary sequence "T-C-A". Complementarity between two single-stranded molecules may be "partial", in which
5 only some of the nucleic acids bind, or it may be complete when total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acids strands.

10 The term "homology", as used herein, refers to a degree of complementarity. There may be partial homology or complete homology (i.e., identity). A partially complementary sequence is one that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid; it is referred to using the functional term "substantially homologous." The inhibition of hybridization of the completely complementary sequence to the target
15 sequence may be examined using a hybridization assay (Southern or northern blot, solution hybridization and the like) under conditions of low stringency. A substantially homologous sequence or probe will compete for and inhibit the binding (i.e., the hybridization) of a completely homologous sequence or probe to the target sequence under conditions of low stringency. This is not to say that conditions of low stringency are such that non-specific
20 binding is permitted; low stringency conditions require that the binding of two sequences to one another be a specific (i.e., selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% identity); in the absence of non-specific binding, the probe will not hybridize to the second non-complementary target sequence.

25 As known in the art, numerous equivalent conditions may be employed to comprise either low or high stringency conditions. Factors such as the length and nature (DNA, RNA, base composition) of the sequence, nature of the target (DNA, RNA, base composition, presence in solution or immobilization, etc.), and the concentration of the salts and other components (e.g., the presence or absence of formamide, dextran sulfate and/or polyethylene
30 glycol) are considered and the hybridization solution may be varied to generate conditions of

either low or high stringency different from, but equivalent to, the above listed conditions.

The term "stringent conditions", as used herein, is the "stringency" which occurs within a range from about $T_m - 5^\circ\text{C}$ (5°C below the melting temperature (T_m) of the probe) to about 20°C to 25°C below T_m . As will be understood by those of skill in the art, the stringency of hybridization may be altered in order to identify or detect identical or related polynucleotide sequences.

The term "antisense", as used herein, refers to nucleotide sequences which are complementary to a specific DNA or RNA sequence. The term "antisense strand" is used in reference to a nucleic acid strand that is complementary to the "sense" strand. Antisense molecules may be produced by any method, including synthesis by ligating the gene(s) of interest in a reverse orientation to a viral promoter which permits the synthesis of a complementary strand. Once introduced into a cell, this transcribed strand combines with natural sequences produced by the cell to form duplexes. These duplexes then block either the further transcription or translation. In this manner, mutant phenotypes may be generated. The designation "negative" is sometimes used in reference to the antisense strand, and "positive" is sometimes used in reference to the sense strand.

The term "portion", as used herein, with regard to a protein (as in "a portion of a given protein") refers to fragments of that protein. The fragments may range in size from four amino acid residues to the entire amino acid sequence minus one amino acid. Thus, a protein "comprising at least a portion of the amino acid sequence of SEQ ID NO:1" encompasses the full-length human HSQEP and fragments thereof.

"Transformation", as defined herein, describes a process by which exogenous DNA enters and changes a recipient cell. It may occur under natural or artificial conditions using various methods well known in the art. Transformation may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method is selected based on the host cell being transformed and may include, but is not limited to, viral infection, electroporation, lipofection, and particle bombardment. Such "transformed" cells include stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome. They also include cells which transiently express the inserted DNA or RNA for limited

periods of time.

The term "antigenic determinant", as used herein, refers to that portion of a molecule that makes contact with a particular antibody (i.e., an epitope). When a protein or fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to a given region or three-dimensional structure on the protein; these regions or structures are referred to as antigenic determinants. An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The terms "specific binding" or "specifically binding", as used herein, in reference to the interaction of an antibody and a protein or peptide, mean that the interaction is dependent upon the presence of a particular structure (i.e., the antigenic determinant or epitope) on the protein; in other words, the antibody is recognizing and binding to a specific protein structure rather than to proteins in general. For example, if an antibody is specific for epitope "A", the presence of a protein containing epitope A (or free, unlabeled A) in a reaction containing labeled "A" and the antibody will reduce the amount of labeled A bound to the antibody.

The term "sample", as used herein, is used in its broadest sense. A biological sample suspected of containing nucleic acid encoding HSQEP or fragments thereof may comprise a cell, chromosomes isolated from a cell (e.g., a spread of metaphase chromosomes), genomic DNA (in solution or bound to a solid support such as for Southern analysis), RNA (in solution or bound to a solid support such as for northern analysis), cDNA (in solution or bound to a solid support), an extract from cells or a tissue, and the like.

The term "correlates with expression of a polynucleotide", as used herein, indicates that the detection of the presence of ribonucleic acid that is similar to SEQ ID NO:2 by northern analysis is indicative of the presence of mRNA encoding HSQEP in a sample and thereby correlates with expression of the transcript from the polynucleotide encoding the protein.

"Alterations" in the polynucleotide of SEQ ID NO: 2, as used herein, comprise any alteration in the sequence of polynucleotides encoding HSQEP including deletions, insertions, and point mutations that may be detected using hybridization assays. Included within this definition is the detection of alterations to the genomic DNA sequence which

encodes HSQEP (e.g., by alterations in the pattern of restriction fragment length polymorphisms capable of hybridizing to SEQ ID NO:2), the inability of a selected fragment of SEQ ID NO: 2 to hybridize to a sample of genomic DNA (e.g., using allele-specific oligonucleotide probes), and improper or unexpected hybridization, such as hybridization to a locus other than the normal chromosomal locus for the polynucleotide sequence encoding HSQEP (e.g., using fluorescence in situ hybridization (FISH) to metaphase chromosomes spreads).

As used herein, the term "antibody" refers to intact molecules as well as fragments thereof, such as Fa, F(ab')₂, and Fv, which are capable of binding the epitopic determinant.

Antibodies that bind HSQEP polypeptides can be prepared using intact polypeptides or fragments containing small peptides of interest as the immunizing antigen. The polypeptide or peptide used to immunize an animal can be derived from translated RNA or synthesized chemically, and can be conjugated to a carrier protein, if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin and thyroglobulin.

The coupled peptide is then used to immunize the animal (e.g., a mouse, a rat, or a rabbit).

The term "humanized antibody", as used herein, refers to antibody molecules in which amino acids have been replaced in the non-antigen binding regions in order to more closely resemble a human antibody, while still retaining the original binding ability.

THE INVENTION

The invention is based on the discovery of a novel human squalene epoxidase (HSQEP), the polynucleotides encoding HSQEP, and the use of these compositions for the diagnosis, prevention, or treatment of hypercholesterolemia and atherosclerosis.

Nucleic acids encoding the human HSQEP of the present invention were first identified in Incyte Clone 638884 from a cDNA library prepared from noncancerous breast tissue removed from a 54-year-old female during a bilateral radical mastectomy (BRSTNOT03) through a computer-generated search for amino acid sequence alignments. A consensus sequence, SEQ ID NO:2, was derived from the following overlapping and/or extended nucleic acid sequences: Incyte Clones 1338208 (COLNNOT13); 000594 (U937NOT01); 013083 (THP1PLB01); 1438629 (PANCNOT08); 043115 (TBLYNOT01); 1434076 (BEPINON01); 1440965 (THYRNOT03); 638884 (BRSTNOT03); and 790619

(PROSTUT03).

In one embodiment, the invention encompasses the novel human squalene epoxidase, a polypeptide comprising the amino acid sequence of SEQ ID NO:1, as shown in Figures 1A, 1B, and 1C. HSQEP is 572 amino acids in length. HSQEP has chemical and structural
5 homology with mouse squalene epoxidase (GI 1217593; SEQ ID NO:3) and rat squalene epoxidase (GI 1083804; SEQ ID NO:4). In particular, HSQEP and mouse squalene epoxidase share 79% identity, while HSQEP and rat squalene epoxidase share 78% identity (Figures 2A, 2B and 2C). As illustrated by Figures 3 and 4, HSQEP and mouse squalene epoxidase have similar hydrophobicity plots. From its amino acid sequence homology with
10 the mouse and rat homologs, HSQEP is predicted to contain a transmembrane domain extending from I₂₁ to V₄₁ of SEQ ID NO:1. In addition, HSQEP contains a putative FAD binding domain extending from I₁₂₆ to E₁₅₂ of SEQ ID NO:1. HSQEP catalyzes the oxidation of squalene to squalene 2,3-epoxide (Figure 5) in the presence of oxygen, FAD, and NADPH.

Many enzymes of the cholesterol biosynthetic pathway are regulated at the
15 transcriptional level. In particular, transcription of these enzymes tends to be up-regulated in the absence of cholesterol in the bloodstream, due to a low-fat diet or treatment with cholesterol-lowering therapeutics. Northern analysis shows that HSQEP transcription appears to follow that trend. Sequences encoding full-length HSQEP were found in thyroid tumor tissue (THYRNOT03) obtained from a donor with a past history of
20 hypercholesterolemia. The donor may have decreased his dietary fat intake and/or have been taking cholesterol-lowering drugs. Sequences encoding full-length HSQEP were also found in healthy colon tissue from a donor with ulcerative colitis. The donor was being treated with anabolic steroids, which may up-regulate transcription of HSQEP. Sequences encoding full-length HSQEP were also found in inflamed adenoid tissue from a three-year old child, a
25 bronchial epithelium primary cell line, and hybrid T-B lymphocytes from a leukemic cell line. It must be noted that expression of HSQEP is not necessarily limited to these tissues.

The invention also encompasses HSQEP variants. A preferred HSQEP variant is one having at least 80%, and more preferably 90%, amino acid sequence similarity to the HSQEP amino acid sequence (SEQ ID NO:1). A most preferred HSQEP variant is one having at least
30 95% amino acid sequence similarity to SEQ ID NO:1.

The invention also encompasses polynucleotides which encode HSQEP. Accordingly, any nucleic acid sequence which encodes the amino acid sequence of HSQEP can be used to generate recombinant molecules which express HSQEP. In a particular embodiment, the invention encompasses the polynucleotide comprising the nucleic acid of
5 SEQ ID NO:2 as shown in Figures 1A, 1B, 1C, 1D, 1E, 1F and 1G.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of nucleotide sequences encoding HSQEP, some bearing minimal homology to the nucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of
10 nucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the nucleotide sequence of naturally occurring HSQEP, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode HSQEP and its variants are preferably
15 capable of hybridizing to the nucleotide sequence of the naturally occurring HSQEP under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding HSQEP or its derivatives possessing a substantially different codon usage. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic expression host in accordance with the frequency
20 with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding HSQEP and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

25 The invention also encompasses production of a DNA sequence, or portions thereof, which encode HSQEP and its derivatives, entirely by synthetic chemistry. After production, the synthetic gene may be inserted into any of the many available DNA vectors and cell systems using reagents that are well known in the art at the time of the filing of this application. Moreover, synthetic chemistry may be used to introduce mutations into a
30 sequence encoding HSQEP or any portion thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed nucleotide sequences, and in particular, those shown in SEQ ID NO:2, under various conditions of stringency. Hybridization conditions are based on the melting temperature (T_m) of the nucleic acid binding complex or probe, as taught in Kimmel, A.R. (1987; Methods Enzymol. Vol. 152), and may be used at a defined stringency.

Altered nucleic acid sequences encoding HSQEP which are encompassed by the invention include deletions, insertions, or substitutions of different nucleotides resulting in a polynucleotide that encodes the same or a functionally equivalent HSQEP. The encoded protein may also contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent HSQEP. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues as long as the biological activity of HSQEP is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid; positively charged amino acids may include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values may include leucine, isoleucine, and valine; glycine and alanine; asparagine and glutamine; serine and threonine; phenylalanine and tyrosine.

Also included within the scope of the present invention are alleles of the gene encoding HSQEP. As used herein, an "allele" or "allelic sequence" is an alternative form of the gene which may result from at least one mutation in the nucleic acid sequence. Alleles may result in altered mRNAs or polypeptides whose structure or function may or may not be altered. Any given gene may have none, one, or many allelic forms. Common mutational changes which give rise to alleles are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

Methods for DNA sequencing which are well known and generally available in the art may be used to practice any embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, Sequenase® (US Biochemical Corp, Cleveland, OH), Taq polymerase (Perkin Elmer), thermostable T7 polymerase (Amersham, Chicago, IL), or combinations of recombinant polymerases and proofreading exonucleases

such as the ELONGASE Amplification System marketed by Gibco BRL (Gaithersburg, MD). Preferably, the process is automated with machines such as the Hamilton Micro Lab 2200 (Hamilton, Reno, NV), Peltier Thermal Cycler (PTC200; MJ Research, Watertown, MA) and the ABI 377 DNA sequencers (Perkin Elmer).

5 The polynucleotide sequence encoding HSQEP may be extended utilizing a partial nucleotide sequence and employing various methods known in the art to detect upstream sequences such as promoters and regulatory elements. For example, one method which may be employed, "restriction-site" PCR, uses universal primers to retrieve unknown sequence adjacent to a known locus (Sarkar, G. (1993) PCR Methods Applic. 2:318-322). In
10 particular, genomic DNA is first amplified in the presence of primer to linker sequence and a primer specific to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

15 Inverse PCR may also be used to amplify or extend sequences using divergent primers based on a known region (Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186). The primers may be designed using OLIGO® 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, MN), or another appropriate program, to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about
20 68°-72° C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

 Another method which may be used is capture PCR which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial
25 chromosome DNA (Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119). In this method, multiple restriction enzyme digestions and ligations may also be used to place an engineered double-stranded sequence into an unknown portion of the DNA molecule before performing PCR.

 Another method which may be used to retrieve unknown sequences is that of Parker,
30 J.D. et al. (1991; Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested

primers, and PromoterFinder™ libraries to walk in genomic DNA (Clontech, Palo Alto, CA). This process avoids the need to screen libraries and may be useful in finding intron/exon junctions.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Also, random-primed libraries are preferable in that they will contain more sequences which contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into the 5' and 3' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity may be converted to electrical signal using appropriate software (e.g. Genotyper™ and Sequence Navigator™, Perkin Elmer) and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode HSQEP, or fusion proteins or functional equivalents thereof, may be used in recombinant DNA molecules to direct expression of HSQEP in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and these sequences may be used to clone and express HSQEP.

As will be understood by those of skill in the art, it may be advantageous to produce HSQEP-encoding nucleotide sequences possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce a recombinant RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated

from the naturally occurring sequence.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter the HSQEP coding sequence for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequence. For example, site-directed mutagenesis may be used to insert new restriction sites, alter glycosylation patterns, to change codon preference, to produce splice variants, or other mutations, and so forth.

In another embodiment of the invention, a natural, modified, or recombinant polynucleotide encoding HSQEP may be ligated to a heterologous sequence to encode a fusion protein. For example, to screen peptide libraries for inhibitors of HSQEP activity, it may be useful to encode a chimeric HSQEP protein that can be recognized by a commercially available antibody. A fusion protein may also be engineered to contain a cleavage site located between a HSQEP encoding sequence and the heterologous protein sequence, so that HSQEP may be cleaved and purified away from the heterologous moiety.

In another embodiment, the coding sequence of HSQEP may be synthesized, in whole or in part, using chemical methods well known in the art (see Caruthers, M.H. et al. (1980) Nucl. Acids Res. Symp. Ser. 215-223, Horn, T. et al. (1980) Nucl. Acids Res. Symp. Ser. 225-232). Alternatively, the protein itself may be produced using chemical methods to synthesize the HSQEP amino acid sequence, or a portion thereof. For example, peptide synthesis can be performed using various solid-phase techniques (Roberge, J.Y. et al. (1995) Science 269:202-204) and automated synthesis may be achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin Elmer).

The newly synthesized peptide may be substantially purified by preparative high performance liquid chromatography (e.g., Creighton, T. (1983) Proteins, Structures and Molecular Principles, WH Freeman and Co., New York, NY). The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure; Creighton, supra). Additionally, the amino acid sequence of HSQEP, or any part thereof, may be altered during direct synthesis and/or combined using chemical

methods with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

In order to express a biologically active HSQEP, the nucleotide sequence encoding HSQEP or functional equivalents, may be inserted into an appropriate expression vector, i.e.,
5 a vector which contains the necessary elements for the transcription and translation of the inserted coding sequence.

Methods which are well known to those skilled in the art may be used to construct expression vectors containing a HSQEP coding sequence and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques,
10 synthetic techniques, and in vivo genetic recombination. Such techniques are described in Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview, NY, and Ausubel, F.M. et al. (1989) Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY.

A variety of expression vector/host systems may be utilized to contain and express a
15 HSQEP coding sequence. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (e.g., baculovirus); plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or
20 with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems.

The "control elements" or "regulatory sequences" are those non-translated regions of the vector--enhancers, promoters, 5' and 3' untranslated regions--which interact with host cellular proteins to carry out transcription, and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of
25 suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the Bluescript® phagemid (Stratagene, LaJolla, CA) or pSport1™ plasmid (Gibco BRL) and ptrp-lac hybrids, and the like may be used. The baculovirus polyhedrin promoter may be used in insect cells. Promoters or
30 enhancers derived from the genomes of plant cells (e.g., heat shock, RUBISCO; and storage

protein genes) or from plant viruses (e.g., viral promoters or leader sequences) may be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding HSQEP, vectors based on SV40 or EBV may be used with an appropriate selectable marker.

In bacterial systems, a number of expression vectors may be selected depending upon the use intended for HSQEP. For example, when large quantities of HSQEP are needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified may be used. Such vectors include, but are not limited to, the

multifunctional *E. coli* cloning and expression vectors such as Bluescript® (Stratagene), in which the sequence encoding HSQEP may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509); and the like. pGEX vectors (Promega, Madison, WI) may also be used to

express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

In the yeast, *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH may be used. For reviews, see Ausubel et al. (supra) and Grant et al. (1987) Methods Enzymol. 153:516-544.

In cases where plant expression vectors are used, the expression of a sequence encoding HSQEP may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV may be used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used (Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105).

These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (see, for example, Hobbs, S. or Murry, L.E. in McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York, NY; pp. 191-196.

5 An insect system may also be used to express HSQEP. For example, in one such system, Autographa californica nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in Spodoptera frugiperda cells or in Trichoplusia larvae. The sequence encoding HSQEP may be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion
10 of HSQEP will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses may then be used to infect, for example, S. frugiperda cells or Trichoplusia larvae in which HSQEP may be expressed (Engelhard, E.K. et al. (1994) Proc. Nat. Acad. Sci. 91:3224-3227).

 In mammalian host cells, a number of viral-based expression systems may be utilized.
15 In cases where an adenovirus is used as an expression vector, a sequence encoding HSQEP may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing HSQEP in infected host cells (Logan, J. and Shenk, T. (1984) Proc. Natl. Acad. Sci. 81:3655-3659). In
20 addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

 Specific initiation signals may also be used to achieve more efficient translation of a sequence encoding HSQEP. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding HSQEP, its initiation codon, and upstream
25 sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a portion thereof, is inserted, exogenous translational control signals including the ATG initiation codon should be provided. Furthermore, the initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational
30 elements and initiation codons may be of various origins, both natural and synthetic. The

efficiency of expression may be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used, such as those described in the literature (Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162).

In addition, a host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to facilitate correct insertion, folding and/or function. Different host cells such as CHO, HeLa, MDCK, HEK293, and WI38, which have specific cellular machinery and characteristic mechanisms for such post-translational activities, may be chosen to ensure the correct modification and processing of the foreign protein.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express HSQEP may be transformed using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or separate vector. Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler, M. et al. (1977) *Cell* 11:223-32) and adenine phosphoribosyltransferase (Lowy, I. et al. (1980) *Cell* 22:817-23) genes which can be employed in *tk⁻* or *aprt⁻* cells, respectively. Also, antimetabolite, antibiotic or herbicide resistance can be used as the basis for selection; for example, *dhfr* which confers resistance to methotrexate (Wigler, M. et al. (1980) *Proc. Natl. Acad. Sci.* 77:3567-70); *npt*, which confers resistance to the aminoglycosides neomycin and G-418 (Colbere-Garapin, F. et al (1981) *J. Mol. Biol.* 150:1-14) and *als* or *pat*, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, *supra*).

Additional selectable genes have been described, for example, *trpB*, which allows cells to utilize indole in place of tryptophan, or *hisD*, which allows cells to utilize histinol in place of histidine (Hartman, S.C. and R.C. Mulligan (1988) *Proc. Natl. Acad. Sci.* 85:8047-51).

Recently, the use of visible markers has gained popularity with such markers as anthocyanins, β glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, being widely used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes, C.A. et al. (1995) *Methods Mol. Biol.* 55:121-131).

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, its presence and expression may need to be confirmed. For example, if the sequence encoding HSQEP is inserted within a marker gene sequence, recombinant cells containing sequences encoding HSQEP can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding HSQEP under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

Alternatively, host cells which contain the coding sequence for HSQEP and express HSQEP may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of the nucleic acid or protein.

The presence of the polynucleotide sequence encoding HSQEP can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes or portions or fragments of polynucleotides encoding HSQEP. Nucleic acid amplification based assays involve the use of oligonucleotides or oligomers based on the HSQEP-encoding sequence to detect transformants containing DNA or RNA encoding HSQEP. As used herein "oligonucleotides" or "oligomers" refer to a nucleic acid sequence of at least about 10 nucleotides and as many as about 60 nucleotides, preferably about 15 to 30 nucleotides, and more preferably about 20-25 nucleotides, which can be used as a probe or amplimer.

A variety of protocols for detecting and measuring the expression of HSQEP, using either polyclonal or monoclonal antibodies specific for the protein are known in the art.

Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on HSQEP is preferred, but a competitive binding assay may be employed. These and other assays are described, among other places, in Hampton, R. et al. (1990; Serological Methods, a Laboratory Manual, APS Press, St Paul, MN) and Maddox, D.E. et al. (1983) J. Exp. Med. 158:1211-1216).

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding HSQEP include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the sequence encoding HSQEP, or any portion of it, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits (Pharmacia & Upjohn, (Kalamazoo, MI); Promega (Madison WI); and U.S. Biochemical Corp., Cleveland, OH). Suitable reporter molecules or labels, which may be used, include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with a nucleotide sequence encoding HSQEP may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a recombinant cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode HSQEP may be designed to contain signal sequences which direct secretion of HSQEP through a prokaryotic or eukaryotic cell membrane. Other recombinant constructions may be used to join sequences encoding HSQEP to nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow

purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, WA). The inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, CA) between the purification domain and HSQEP may be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing HSQEP and a nucleic acid encoding 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification on IMIAC (immobilized metal ion affinity chromatography as described in Porath, J. et al. (1992, Prot. Exp. Purif. 3:263-281) while the enterokinase cleavage site provides a means for purifying HSQEP from the fusion protein. A discussion of vectors which contain fusion proteins is provided in Kroll, D.J. et al. (1993; DNA Cell Biol. 12:441-453).

In addition to recombinant production, fragments of HSQEP may be produced by direct peptide synthesis using solid-phase techniques (cf Stewart et al. (1969) Solid-Phase Peptide Synthesis, W.H. Freeman Co., San Francisco, CA; Merrifield J. (1963) J. Am. Chem. Soc. 85:2149-2154). Protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Various fragments of HSQEP may be chemically synthesized separately and combined using chemical methods to produce the full length molecule

THERAPEUTICS

In another embodiment of the invention, HSQEP or fragments thereof may be used for therapeutic purposes.

As shown in Figure 5, mevalonate, GPP and FPP are not only precursors of cholesterol, but are also the precursors of mevalonate-derived non-sterol metabolites such as ubiquinones, dolichols, and isoprenic acids. Lovastatin is a widely-used cholesterol-lowering drug which inhibits HMG-CoA reductase, an enzyme at the early stage in the cholesterol biosynthetic pathway. Consequently, in addition to cholesterol, lovastatin and other HMG-CoA reductase inhibitors deplete the important mevalonate-derived non-sterol metabolites. There is a significant need for anti-hypercholesterolemic therapeutics which do

not also deplete the production of mevalonate-derived non-sterol metabolites. Since squalene is much further along the pathway of cholesterol biosynthesis, post-FPP and the non-sterol metabolite branch point (Figure 5), HSQEP is a much better target for the design of cholesterol-lowering therapeutics than HMG-CoA reductase.

5 Therefore, in one embodiment, antagonists or inhibitors which block or modulate the activity of HSQEP may be used in those situations where such inhibition is therapeutically desirable. Such antagonists or inhibitors may be produced using a variety of methods which are generally known in the art. In particular, purified HSQEP may be used to screen libraries of pharmaceutical agents for those which specifically bind HSQEP or to produce antibodies.

10 In one aspect, antibodies which are specific for HSQEP may be used directly as an antagonist, or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissue which express HSQEP. The antibodies may be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, Fab fragments, and fragments produced by a
15 Fab expression library. Neutralizing antibodies, (i.e., those which inhibit dimer formation) are especially preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others, may be immunized by injection with HSQEP or any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species,
20 various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

25 It is preferred that the peptides, fragments, or oligopeptides used to induce antibodies to HSQEP have an amino acid sequence consisting of at least five amino acids, and more preferably at least 10 amino acids. It is also preferable that they are identical to a portion of the amino acid sequence of the natural protein, and they may contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of HSQEP amino acids
30 may be fused with those of another protein such as keyhole limpet hemocyanin and antibody

produced against the chimeric molecule.

Monoclonal antibodies to HSQEP may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma
5 technique, and the EBV-hybridoma technique (Koehler et al. (1975) Nature 256:495-497; Kosbor et al. (1983) Immunol. Today 4:72; Cote et al. (1983) Proc. Natl. Acad. Sci. 80:2026-2030; Cole et al. (1985) Monoclonal Antibodies and Cancer Therapy, Alan R. Liss Inc., New York, NY, pp. 77-96).

In addition, techniques developed for the production of "chimeric antibodies", the
10 splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity can be used (Morrison et al. (1984) Proc. Natl. Acad. Sci. 81:6851-6855; Neuberger et al. (1984) Nature 312:604-608; Takeda et al. (1985) Nature 314:452-454). Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce
15 HSQEP-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries (Burton D.R. (1991) Proc. Natl. Acad. Sci. 88:11120-3).

Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening recombinant immunoglobulin libraries or panels of highly specific
20 binding reagents as disclosed in the literature (Orlandi, et al. (1989) Proc. Natl. Acad. Sci. 86: 3833-3837; Winter, G. et al. (1991) Nature 349:293-299).

Antibody fragments which contain specific binding sites for HSQEP may also be generated. For example, such fragments include, but are not limited to, the F(ab')₂ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments
25 which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity (Huse et al. (1989) Science 256:1275-1281).

Various immunoassays may be used for screening to identify antibodies having the
30 desired specificity. Numerous protocols for competitive binding or immunoradiometric

assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between HSQEP and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering HSQEP epitopes is preferred, but a competitive binding assay may also be employed (Maddox, supra).

In another embodiment of the invention, the polynucleotides encoding HSQEP, or any fragment thereof, or antisense sequences, may be used for therapeutic purposes. In one aspect, antisense to the polynucleotide encoding HSQEP may be used in situations in which it would be desirable to block the synthesis of the protein. In particular, cells may be transformed with sequences complementary to polynucleotides encoding HSQEP. Thus, antisense sequences may be used to modulate HSQEP activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligomers or larger fragments, can be designed from various locations along the coding or control regions of sequences encoding HSQEP.

Expression vectors derived from retroviruses, adenovirus, herpes or vaccinia viruses, or from various bacterial plasmids may be used for delivery of nucleotide sequences to the targeted organ, tissue or cell population. Methods which are well known to those skilled in the art can be used to construct recombinant vectors which will express antisense polynucleotides of the gene encoding HSQEP. These techniques are described both in Sambrook et al. (supra) and in Ausubel et al. (supra).

Genes encoding HSQEP can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide or fragment thereof which encodes HSQEP. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until all copies are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector and even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing antisense molecules, DNA, RNA, or PNA, to the control regions of the gene encoding

HSQEP, i.e., the promoters, enhancers, and introns. Oligonucleotides derived from the transcription initiation site, e.g., between positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature (Gee, J.E. et al. (1994) In: Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing Co., Mt. Kisco, NY). The antisense molecules may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Examples which may be used include engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding HSQEP.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target gene containing the cleavage site may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Antisense molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of RNA molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding HSQEP. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as

T7 or SP6. Alternatively, these cDNA constructs that synthesize antisense RNA constitutively or inducibly can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection and by liposome injections may be achieved using methods which are well known in the art.

Any of the therapeutic methods described above may be applied to any suitable subject including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

An additional embodiment of the invention relates to the administration of a pharmaceutical composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of HSQEP, antibodies to HSQEP, mimetics, agonists, antagonists, or inhibitors of HSQEP. The compositions may be administered alone or in combination with at least one other agent, such as stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous,

intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such

as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated
5 in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active
10 compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to
15 be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

20 The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1-50 mM histidine, 0.1%-2% sucrose,
25 and 2-7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of HSQEP, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions
30 wherein the active ingredients are contained in an effective amount to achieve the intended

purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models, usually mice, rabbits, dogs, or pigs. The animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example HSQEP or fragments thereof, antibodies of HSQEP, agonists, antagonists or inhibitors of HSQEP, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., ED50 (the dose therapeutically effective in 50% of the population) and LD50 (the dose lethal to 50% of the population). The dose ratio between therapeutic and toxic effects is the therapeutic index, and it can be expressed as the ratio, LD50/ED50.

Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from 0.1 to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages

and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

5 **DIAGNOSTICS**

In another embodiment, antibodies which specifically bind HSQEP may be used for the diagnosis of conditions or diseases characterized by expression of HSQEP, or in assays to monitor patients being treated with HSQEP, agonists, antagonists or inhibitors. The antibodies useful for diagnostic purposes may be prepared in the same manner as those described above for therapeutics. Diagnostic assays for HSQEP include methods which utilize the antibody and a label to detect HSQEP in human body fluids or extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by joining them, either covalently or non-covalently, with a reporter molecule. A wide variety of reporter molecules which are known in the art may be used, several of which are described above.

A variety of protocols including ELISA, RIA, and FACS for measuring HSQEP are known in the art and provide a basis for diagnosing altered or abnormal levels of HSQEP expression. Normal or standard values for HSQEP expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to HSQEP under conditions suitable for complex formation. The amount of standard complex formation may be quantified by various methods, but preferably by photometric, means. Quantities of HSQEP expressed in subject, control and disease, samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding HSQEP may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, antisense RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantitate gene expression in biopsied tissues in which expression of HSQEP may be correlated with disease. The diagnostic assay may be used to distinguish between absence, presence, and excess expression of HSQEP, and to

monitor regulation of HSQEP levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding HSQEP or closely related molecules, may be used to identify nucleic acid sequences which encode HSQEP. The
5 specificity of the probe, whether it is made from a highly specific region, e.g., 10 unique nucleotides in the 5' regulatory region, or a less specific region, e.g., especially in the 3' coding region, and the stringency of the hybridization or amplification (maximal, high, intermediate, or low) will determine whether the probe identifies only naturally occurring sequences encoding HSQEP, alleles, or related sequences.

10 Probes may also be used for the detection of related sequences, and should preferably contain at least 50% of the nucleotides from any of the HSQEP encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and derived from the nucleotide sequence of SEQ ID NO:2 or from genomic sequence including promoter, enhancer elements, and introns of the naturally occurring HSQEP.

15 Means for producing specific hybridization probes for DNAs encoding HSQEP include the cloning of nucleic acid sequences encoding HSQEP or HSQEP derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes
20 may be labeled by a variety of reporter groups, for example, radionuclides such as ³²P or ³⁵S, or enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding HSQEP may be used for the diagnosis of conditions or diseases which are associated with expression of HSQEP. Examples of such
25 conditions or diseases include disorders of cholesterol metabolism such as hypercholesterolemia. The polynucleotide sequences encoding HSQEP may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; or in dip stick, pin, ELISA or chip assays utilizing fluids or tissues from patient biopsies to detect altered HSQEP expression. Such qualitative or quantitative methods are
30 well known in the art.

In a particular aspect, the nucleotide sequences encoding HSQEP may be useful in assays that detect activation or induction of various cancers, particularly those mentioned above. The nucleotide sequence encoding HSQEP may be labeled by standard methods, and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the biopsied or extracted sample is significantly altered from that of a comparable control sample, the nucleotide sequence has hybridized with nucleotide sequences in the sample, and the presence of altered levels of nucleotide sequences encoding HSQEP in the sample indicates the presence of the associated disease. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or in monitoring the treatment of an individual patient.

In order to provide a basis for the diagnosis of disease associated with expression of HSQEP, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, which encodes HSQEP, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with those from an experiment where a known amount of a substantially purified polynucleotide is used. Standard values obtained from normal samples may be compared with values obtained from samples from patients who are symptomatic for disease. Deviation between standard and subject values is used to establish the presence of disease.

Once disease is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to evaluate whether the level of expression in the patient begins to approximate that which is observed in the normal patient. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual

clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides encoding HSQEP may involve the use
5 of PCR. Such oligomers may be chemically synthesized, generated enzymatically, or produced from a recombinant source. Oligomers will preferably consist of two nucleotide sequences, one with sense orientation (5'→3') and another with antisense (3'←5'), employed under optimized conditions for identification of a specific gene or condition. The same two oligomers, nested sets of oligomers, or even a degenerate pool of oligomers may be employed
10 under less stringent conditions for detection and/or quantitation of closely related DNA or RNA sequences.

Methods which may also be used to quantitate the expression of HSQEP include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and standard curves onto which the experimental results are interpolated (Melby, P.C. et al.
15 (1993) J. Immunol. Methods, 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 229-236). The speed of quantitation of multiple samples may be accelerated by running the assay in an ELISA format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In another embodiment of the invention, the nucleic acid sequence which encodes
20 HSQEP may also be used to generate hybridization probes which are useful for mapping the naturally occurring genomic sequence. The sequence may be mapped to a particular chromosome or to a specific region of the chromosome using well known techniques. Such techniques include FISH, FACS, or artificial chromosome constructions, such as yeast artificial chromosomes, bacterial artificial chromosomes, bacterial P1 constructions or single
25 chromosome cDNA libraries as reviewed in Price, C.M. (1993) Blood Rev. 7:127-134, and Trask, B.J. (1991) Trends Genet. 7:149-154.

FISH (as described in Verma et al. (1988) Human Chromosomes: A Manual of Basic Techniques, Pergamon Press, New York, NY) may be correlated with other physical chromosome mapping techniques and genetic map data. Examples of genetic map data can
30 be found in the 1994 Genome Issue of Science (265:1981f). Correlation between the location

of the gene encoding HSQEP on a physical chromosomal map and a specific disease , or predisposition to a specific disease, may help delimit the region of DNA associated with that genetic disease. The nucleotide sequences of the subject invention may be used to detect differences in gene sequences between normal, carrier, or affected individuals.

5 In situ hybridization of chromosomal preparations and physical mapping techniques such as linkage analysis using established chromosomal markers may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to
10 chromosomal arms, or parts thereof, by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, for example, AT to 11q22-23 (Gatti, R.A. et al. (1988) Nature 336:577-580), any sequences mapping to that area may represent associated or
15 regulatory genes for further investigation. The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc. among normal, carrier, or affected individuals.

 In another embodiment of the invention, HSQEP, its catalytic or immunogenic fragments or oligopeptides thereof, can be used for screening libraries of compounds in any
20 of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes, between HSQEP and the agent being tested, may be measured.

 Another technique for drug screening which may be used provides for high
25 throughput screening of compounds having suitable binding affinity to the protein of interest as described in published PCT application WO84/03564. In this method, as applied to HSQEP large numbers of different small test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with HSQEP, or fragments thereof, and washed. Bound HSQEP is then detected by methods well known in
30 the art. Purified HSQEP can also be coated directly onto plates for use in the aforementioned

drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding HSQEP specifically compete with a test compound
5 for binding HSQEP. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with HSQEP.

In additional embodiments, the nucleotide sequences which encode HSQEP may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but
10 not limited to, such properties as the triplet genetic code and specific base pair interactions.

The examples below are provided to illustrate the subject invention and are not included for the purpose of limiting the invention.

EXAMPLES

15 I cDNA Library Construction

The BRSTNOT03 cDNA library was constructed from tissue removed from the normal breast of a 54 year old female (Lot No. 0025B; Mayo Clinic, Rochester, MN). The frozen tissue was immediately homogenized and lysed using a Brinkmann Homogenizer Polytron PT-3000 (Brinkmann Instruments Inc, Westbury NY) in guanidinium isothiocyanate
20 solution. Lysates were then loaded on a 5.7 M CsCl cushion and ultracentrifuged in a SW28 swinging bucket rotor for 18 hours at 25,000 rpm at ambient temperature. The RNA was extracted once with acid phenol at pH 4.0 and once with phenol chloroform at pH 8.0 and precipitated using 0.3 M sodium acetate and 2.5 volumes of ethanol, resuspended in DEPC-treated water and DNase treated for 25 min at 37°C. The reaction was stopped with
25 an equal volume of acid phenol, and the RNA was isolated using the Qiagen Oligotex kit (Qiagen Inc., Chatsworth, CA) and used to construct the cDNA library.

The RNA was handled according to the recommended protocols in the SuperScript Plasmid System for cDNA Synthesis and Plasmid Cloning (Cat. No. 18248-013; Gibco/BRL). cDNAs were fractionated on a Sepharose CL4B column (Cat. No. 275105, Pharmacia), and those cDNAs exceeding 400 bp were ligated into pSport I. The plasmid
30

pSport I was subsequently transformed into DH5a™ competent cells (Cat. No. 18258-012, Gibco/BRL).

II Isolation and Sequencing of cDNA Clones

Plasmid DNA was released from the cells and purified using the Miniprep Kit (Cat. No. 77468; Advanced Genetic Technologies Corporation, Gaithersburg MD). This kit consists of a 96 well block with reagents for 960 purifications. The recommended protocol was employed except for the following changes: 1) the 96 wells were each filled with only 1 ml of sterile Terrific Broth (Cat. No. 22711, Life Technologies, Gaithersburg, MD) with carbenicillin at 25 mg/L and glycerol at 0.4%; 2) the bacteria were cultured for 24 hours after the wells were inoculated and then lysed with 60 µl of lysis buffer; 3) a centrifugation step employing the Beckman GS-6R @2900 rpm for 5 min was performed before the contents of the block were added to the primary filter plate; and 4) the optional step of adding isopropanol to TRIS buffer was not routinely performed. After the last step in the protocol, samples were transferred to a Beckman 96-well block for storage.

The cDNAs were sequenced by the method of Sanger F. and A.R. Coulson (1975; J. Mol. Biol. 94:441f), using a Hamilton Micro Lab 2200 (Hamilton, Reno NV) in combination with four Peltier Thermal Cyclers (PTC200; MJ Research, Watertown MA) and Applied Biosystems 377 or 373 DNA Sequencing Systems (Perkin Elmer), and reading frame was determined.

III Homology Searching of cDNA Clones and Their Deduced Proteins

Each cDNA was compared to sequences in GenBank using a search algorithm developed by Applied Biosystems and incorporated into the INHERIT™ 670 sequence analysis system. In this algorithm, Pattern Specification Language (TRW Inc, Los Angeles, CA) was used to determine regions of homology. The three parameters that determine how the sequence comparisons run were window size, window offset, and error tolerance. Using a combination of these three parameters, the DNA database was searched for sequences containing regions of homology to the query sequence, and the appropriate sequences were scored with an initial value. Subsequently, these homologous regions were examined using dot matrix homology plots to distinguish regions of homology from chance matches. Smith-Waterman alignments were used to display the results of the homology search.

Peptide and protein sequence homologies were ascertained using the INHERIT- 670 sequence analysis system using the methods similar to those used in DNA sequence homologies. Pattern Specification Language and parameter windows were used to search protein databases for sequences containing regions of homology which were scored with an
5 initial value. Dot-matrix homology plots were examined to distinguish regions of significant homology from chance matches.

BLAST, which stands for Basic Local Alignment Search Tool (Altschul, S.F. (1993) J. Mol. Evol. 36:290-300; Altschul et al. (1990) J. Mol. Biol. 215:403-410), was used to search for local sequence alignments. BLAST produces alignments of both nucleotide and
10 amino acid sequences to determine sequence similarity. Because of the local nature of the alignments, BLAST is especially useful in determining exact matches or in identifying homologs. BLAST is useful for matches which do not contain gaps. The fundamental unit of BLAST algorithm output is the High-scoring Segment Pair (HSP).

An HSP consists of two sequence fragments of arbitrary but equal lengths whose
15 alignment is locally maximal and for which the alignment score meets or exceeds a threshold or cutoff score set by the user. The BLAST approach is to look for HSPs between a query sequence and a database sequence, to evaluate the statistical significance of any matches found, and to report only those matches which satisfy the user-selected threshold of significance. The parameter E establishes the statistically significant threshold for reporting
20 database sequence matches. E is interpreted as the upper bound of the expected frequency of chance occurrence of an HSP (or set of HSPs) within the context of the entire database search. Any database sequence whose match satisfies E is reported in the program output.

IV Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript
25 of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound (Sambrook et al., supra).

Analogous computer techniques using BLAST (Altschul, S.F. 1993 and 1990, supra) are used to search for identical or related molecules in nucleotide databases such as GenBank or the LIFESEQ™ database (Incyte Pharmaceuticals). This analysis is much faster than
30 multiple, membrane-based hybridizations. In addition, the sensitivity of the computer search

can be modified to determine whether any particular match is categorized as exact or homologous.

The basis of the search is the product score which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1-2% error; and at 70, the match will be exact. Homologous molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

The results of northern analysis are reported as a list of libraries in which the transcript encoding HSQEP occurs. Abundance and percent abundance are also reported. Abundance directly reflects the number of times a particular transcript is represented in a cDNA library, and percent abundance is abundance divided by the total number of sequences examined in the cDNA library.

V Extension of HSQEP-Encoding Polynucleotides to Full Length or to Recover Regulatory Sequences

Full length HSQEP-encoding nucleic acid sequence (SEQ ID NO:2) is used to design oligonucleotide primers for extending a partial nucleotide sequence to full length or for obtaining 5' or 3', intron or other control sequences from genomic libraries. One primer is synthesized to initiate extension in the antisense direction (XLR) and the other is synthesized to extend sequence in the sense direction (XLF). Primers are used to facilitate the extension of the known sequence "outward" generating amplicons containing new, unknown nucleotide sequence for the region of interest. The initial primers are designed from the cDNA using OLIGO 4.06 (National Biosciences), or another appropriate program, to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68°-72° C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations is avoided.

The original, selected cDNA libraries, or a human genomic library are used to extend the sequence; the latter is most useful to obtain 5' upstream regions. If more extension is

necessary or desired, additional sets of primers are designed to further extend the known region.

By following the instructions for the XL-PCR kit (Perkin Elmer) and thoroughly mixing the enzyme and reaction mix, high fidelity amplification is obtained. Beginning with
 5 40 pmol of each primer and the recommended concentrations of all other components of the kit, PCR is performed using the Peltier Thermal Cycler (PTC200; M.J. Research, Watertown, MA) and the following parameters:

	Step 1	94° C for 1 min (initial denaturation)
	Step 2	65° C for 1 min
10	Step 3	68° C for 6 min
	Step 4	94° C for 15 sec
	Step 5	65° C for 1 min
	Step 6	68° C for 7 min
	Step 7	Repeat step 4-6 for 15 additional cycles
15	Step 8	94° C for 15 sec
	Step 9	65° C for 1 min
	Step 10	68° C for 7:15 min
	Step 11	Repeat step 8-10 for 12 cycles
	Step 12	72° C for 8 min
20	Step 13	4° C (and holding)

A 5-10 μ l aliquot of the reaction mixture is analyzed by electrophoresis on a low concentration (about 0.6-0.8%) agarose mini-gel to determine which reactions were successful in extending the sequence. Bands thought to contain the largest products are
 25 selected and removed from the gel. Further purification involves using a commercial gel extraction method such as QIAQuick™ (QIAGEN Inc., Chatsworth, CA). After recovery of the DNA, Klenow enzyme is used to trim single-stranded, nucleotide overhangs creating blunt ends which facilitate religation and cloning.

After ethanol precipitation, the products are redissolved in 13 μ l of ligation buffer,
 30 1 μ l T4-DNA ligase (15 units) and 1 μ l T4 polynucleotide kinase are added, and the mixture is incubated at room temperature for 2-3 hours or overnight at 16° C. Competent *E. coli* cells (in 40 μ l of appropriate media) are transformed with 3 μ l of ligation mixture and cultured in 80 μ l of SOC medium (Sambrook et al., supra). After incubation for one hour at 37° C, the whole transformation mixture is plated on Luria Bertani (LB)-agar (Sambrook et al., supra)

containing 2x Carb. The following day, several colonies are randomly picked from each plate and cultured in 150 μ l of liquid LB/2x Carb medium placed in an individual well of an appropriate, commercially-available, sterile 96-well microtiter plate. The following day, 5 μ l of each overnight culture is transferred into a non-sterile 96-well plate and after dilution 1:10 with water, 5 μ l of each sample is transferred into a PCR array.

For PCR amplification, 18 μ l of concentrated PCR reaction mix (3.3x) containing 4 units of rTth DNA polymerase, a vector primer, and one or both of the gene specific primers used for the extension reaction are added to each well. Amplification is performed using the following conditions:

10	Step 1	94° C for 60 sec
	Step 2	94° C for 20 sec
	Step 3	55° C for 30 sec
	Step 4	72° C for 90 sec
	Step 5	Repeat steps 2-4 for an additional 29 cycles
15	Step 6	72° C for 180 sec
	Step 7	4° C (and holding)

Aliquots of the PCR reactions are run on agarose gels together with molecular weight markers. The sizes of the PCR products are compared to the original partial cDNAs, and appropriate clones are selected, ligated into plasmid, and sequenced.

20 VI Labeling and Use of Hybridization Probes

Hybridization probes derived from SEQ ID NO:2 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base-pairs, is specifically described, essentially the same procedure is used with larger cDNA fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 (National Biosciences), labeled by combining 50 pmol of each oligomer and 250 mCi of [γ -³²P] adenosine triphosphate (Amersham) and T4 polynucleotide kinase (DuPont NEN®, Boston, MA). The labeled oligonucleotides are substantially purified with Sephadex G-25 superfine resin column (Pharmacia & Upjohn). A portion containing 10⁷ counts per minute of each of the sense and antisense oligonucleotides is used in a typical membrane based hybridization analysis of human genomic DNA digested with one of the following endonucleases (Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II; DuPont NEN®).

The DNA from each digest is fractionated on a 0.7 percent agarose gel and

transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham, NH).

Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under increasingly stringent conditions up to 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. After XOMAT AR™ film (Kodak, Rochester, NY) is exposed to the blots in a Phosphorimager cassette (Molecular Dynamics, Sunnyvale, CA) for several hours, hybridization patterns are compared visually.

VII Antisense Molecules

Antisense molecules to the HSQEP-encoding sequence, or any part thereof, is used to inhibit in vivo or in vitro expression of naturally occurring HSQEP. Although use of antisense oligonucleotides, comprising about 20 base-pairs, is specifically described, essentially the same procedure is used with larger cDNA fragments. An oligonucleotide based on the coding sequences of HSQEP, as shown in Figures 1A, 1B, 1C, 1D, 1E, 1F and 1G, is used to inhibit expression of naturally occurring HSQEP. The complementary oligonucleotide is designed from the most unique 5' sequence as shown in Figures 1A, 1B, 1C, 1D, 1E, 1F and 1G and used either to inhibit transcription by preventing promoter binding to the upstream nontranslated sequence or translation of an HSQEP-encoding transcript by preventing the ribosome from binding. Using an appropriate portion of the signal and 5' sequence of SEQ ID NO:2, an effective antisense oligonucleotide includes any 15-20 nucleotides spanning the region which translates into the signal or 5' coding sequence of the polypeptide as shown in Figures 1A, 1B, 1C, 1D, 1E, 1F and 1G.

VIII Expression of HSQEP

Expression of HSQEP is accomplished by subcloning the cDNAs into appropriate vectors and transforming the vectors into host cells. In this case, the cloning vector, pSport, previously used for the generation of the cDNA library is used to express HSQEP in E. coli. Upstream of the cloning site, this vector contains a promoter for β -galactosidase, followed by sequence containing the amino-terminal Met, and the subsequent seven residues of β -galactosidase. Immediately following these eight residues is a bacteriophage promoter useful for transcription and a linker containing a number of unique restriction sites.

Induction of an isolated, transformed bacterial strain with IPTG using standard methods produces a fusion protein which consists of the first eight residues of

β -galactosidase, about 5 to 15 residues of linker, and the full length protein. The signal residues direct the secretion of HSQEP into the bacterial growth media which can be used directly in the following assay for activity.

IX Demonstration of HSQEP Activity

5 The HSQEP-catalyzed oxidation of squalene to squalene 2,3-epoxide is assayed as described for rat squalene epoxidase (Sakakibara, J. et al. (1995) J. Biol. Chem. 270:17-20). Assay mixtures contain HSQEP, 20 mM Tris-HCL pH 7.5, 0.01 mM FAD, 0.2 units of NADPH-cytochrome C (P-450) reductase, 0.01 mM [14 C]-squalene (dispersed with the aid of 20 ml Tween-80), 0.2% Triton X-100, and 1 mM NADPH in a total volume of 0.5 ml. The
10 reaction is initiated by addition of NADPH. Reaction mixtures are incubated at 37°C for 30 minutes. Reaction products and [14 C]-lipid standards are analyzed by silica gel thin-layer chromatography developed in a 0.5/95.5 (v/v) mixture of ethyl acetate/benzene, followed by autoradiography.

X Production of HSQEP Specific Antibodies

15 HSQEP that is substantially purified using PAGE electrophoresis (Sambrook, supra), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols. The amino acid sequence deduced from SEQ ID NO:2 is analyzed using DNASTAR software (DNASTAR Inc) to determine regions of high immunogenicity and a corresponding oligopeptide is synthesized and used to raise antibodies by means known
20 to those of skill in the art. Selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions, is described by Ausubel et al. (supra), and others.

Typically, the oligopeptides are 15 residues in length, synthesized using an Applied Biosystems Peptide Synthesizer Model 431A using fmoc-chemistry, and coupled to keyhole limpet hemocyanin (KLH, Sigma, St. Louis, MO) by reaction with N-maleimidobenzoyl-N-
25 hydroxysuccinimide ester (MBS; Ausubel et al., supra). Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. The resulting antisera are tested for antipeptide activity, for example, by binding the peptide to plastic, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radioiodinated, goat anti-rabbit IgG.

XI Purification of Naturally Occurring HSQEP Using Specific Antibodies

Naturally occurring or recombinant HSQEP is substantially purified by immunoaffinity chromatography using antibodies specific for HSQEP. An immunoaffinity column is constructed by covalently coupling HSQEP antibody to an activated
5 chromatographic resin, such as CnBr-activated Sepharose (Pharmacia & Upjohn). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing HSQEP is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of HSQEP (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that
10 disrupt antibody/HSQEP binding (eg, a buffer of pH 2-3 or a high concentration of a chaotrope, such as urea or thiocyanate ion), and HSQEP is collected.

XII Identification of Molecules Which Interact with HSQEP

HSQEP or biologically active fragments thereof are labeled with ¹²⁵I Bolton-Hunter reagent (Bolton et al. (1973) Biochem. J. 133: 529). Candidate molecules previously arrayed
15 in the wells of a multi-well plate are incubated with the labeled HSQEP, washed and any wells with labeled HSQEP complex are assayed. Data obtained using different concentrations of HSQEP are used to calculate values for the number, affinity, and association of HSQEP with the candidate molecules.

All publications and patents mentioned in the above specification are herein
20 incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of
25 the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION

- (i) APPLICANT: INCYTE PHARMACEUTICALS, INC.
- (ii) TITLE OF THE INVENTION: HUMAN SQUALENE EPOXIDASE
- (iii) NUMBER OF SEQUENCES: 4
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Incyte Pharmaceuticals, Inc.
 - (B) STREET: 3174 Porter Drive
 - (C) CITY: Palo Alto
 - (D) STATE: California
 - (E) COUNTRY: USA
 - (F) ZIP: 94304
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Diskette
 - (B) COMPUTER: IBM Compatible
 - (C) OPERATING SYSTEM: DOS
 - (D) SOFTWARE: FastSEQ Version 1.5
- (vi) CURRENT APPLICATION DATA:
 - (A) PCT APPLICATION NUMBER: To Be Assigned
 - (B) FILING DATE: Herewith
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 08/745,934
 - (B) FILING DATE: 07-NOV-1996
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Billings, Lucy J.
 - (B) REGISTRATION NUMBER: 36,749
 - (C) REFERENCE/DOCKET NUMBER: PF-0151 PCT
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (650) 855-0555
 - (B) TELEFAX: (650) 845-4166

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 572 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: peptide
- (vii) IMMEDIATE SOURCE:
 - (A) LIBRARY:
 - (B) CLONE: Consensus
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

Met Trp Thr Phe Leu Gly Ile Ala Thr Phe Thr Tyr Phe Tyr Lys Lys
 1           5           10           15
Phe Gly Asp Phe Ile Thr Leu Ala Asn Arg Glu Val Leu Leu Cys Val
 20           25           30
Leu Val Phe Leu Ser Leu Gly Leu Val Leu Ser Tyr Arg Cys Arg His
 35           40           45
Arg Asn Gly Gly Leu Leu Gly Arg Gln Lys Ser Gly Ser Gln Ile Ala
 50           55           60

```

Leu	Phe	Ser	Asp	Ile	Leu	Ser	Gly	Leu	Pro	Phe	Ile	Gly	Phe	Phe	Trp
65					70					75					80
Ala	Asn	Pro	Pro	Leu	Asn	Gln	Lys	Ile	Arg	Ser	Ser	Ser	Arg	Gln	Glu
				85					90					95	
Ala	Gln	Lys	Arg	Asn	Gln	Tyr	Phe	Arg	Asn	Lys	Leu	Asn	Arg	Asn	Ser
			100					105					110		
Cys	Cys	Thr	Ser	Thr	Ser	Ser	Gln	Asn	Asp	Pro	Glu	Val	Ile	Ile	Val
		115					120					125			
Gly	Ala	Gly	Val	Leu	Gly	Ser	Ala	Leu	Ala	Ala	Val	Leu	Ser	Arg	Asp
130						135					140				
Gly	Arg	Lys	Val	Thr	Val	Ile	Glu	Arg	Asp	Leu	Lys	Glu	Pro	Asp	Arg
145					150					155					160
Ile	Val	Gly	Glu	Phe	Leu	Gln	Pro	Gly	Gly	Tyr	His	Val	Leu	Lys	Asp
				165				170						175	
Leu	Gly	Leu	Gly	Asp	Thr	Val	Glu	Gly	Leu	Asp	Ala	Gln	Val	Val	Asn
			180					185					190		
Gly	Tyr	Met	Ile	His	Asp	Gln	Glu	Ser	Lys	Ser	Glu	Val	Gln	Ile	Pro
		195					200					205			
Asn	Pro	Leu	Ser	Glu	Asn	Asn	Gln	Val	Gln	Ser	Gly	Arg	Ala	Phe	His
210						215						220			
His	Gly	Arg	Phe	Ile	Met	Ser	Leu	Arg	Lys	Ala	Val	Met	Ala	Glu	Pro
225					230					235					240
Asn	Ala	Lys	Phe	Ile	Glu	Gly	Val	Val	Leu	Gln	Leu	Leu	Glu	Glu	Asp
				245					250					255	
Asp	Val	Val	Met	Gly	Val	Gln	His	Lys	Asp	Lys	Glu	Thr	Gly	Asp	Ile
			260					265					270		
Lys	Glu	Leu	His	Ala	Pro	Leu	Thr	Val	Val	Ala	Asp	Gly	Leu	Phe	Ser
			275				280					285			
Lys	Phe	Arg	Lys	Ser	Leu	Val	Ser	Asn	Lys	Val	Ser	Val	Ser	Ser	His
290						295					300				
Phe	Val	Gly	Phe	Leu	Met	Lys	Asn	Ala	Pro	Gln	Phe	Thr	Ala	Asn	His
305					310					315					320
Ala	Glu	Leu	Ile	Leu	Ala	Asn	Pro	Ser	Pro	Val	Leu	Ile	Tyr	Gln	Ile
				325					330					335	
Ser	Ser	Ser	Glu	Ile	Glu	Tyr	Leu	Leu	Thr	Leu	Glu	Gly	Met	Pro	Arg
			340				345						350		
Asn	Leu	Arg	Glu	Tyr	Met	Val	Glu	Lys	Ile	Tyr	Pro	Gln	Ile	Pro	Asp
			355				360					365			
His	Leu	Lys	Glu	Pro	Phe	Leu	Glu	Ala	Thr	Asp	Asn	Ser	His	Leu	Arg
			370			375					380				
Ser	Met	Pro	Ala	Ser	Phe	Leu	Pro	Pro	Ser	Ser	Val	Lys	Lys	Arg	Gly
385					390					395					400
Val	Leu	Leu	Leu	Gly	Asp	Ala	Tyr	Asn	Met	Arg	His	Pro	Leu	Thr	Gly
				405					410					415	
Gly	Gly	Met	Thr	Val	Ala	Phe	Lys	Asp	Ile	Lys	Leu	Trp	Arg	Lys	Leu
			420					425					430		
Leu	Lys	Gly	Ile	Pro	Asp	Leu	Tyr	Asp	Asp	Ala	Ala	Ile	Phe	Glu	Ala
		435					440					445			
Lys	Lys	Ser	Phe	Tyr	Trp	Ala	Arg	Lys	Thr	Ser	His	Ser	Phe	Val	Val
450						455					460				
Asn	Ile	Leu	Ala	Gln	Ala	Leu	Tyr	Glu	Leu	Phe	Ser	Ala	Thr	Asp	Asp
465					470					475					480
Ser	Leu	His	Gln	Leu	Arg	Lys	Ala	Cys	Phe	Leu	Tyr	Phe	Lys	Leu	Gly
				485					490					495	
Gly	Glu	Cys	Xaa	Ala	Gly	Pro	Val	Gly	Leu	Leu	Ser	Val	Leu	Ser	Pro
			500					505					510		
Asn	Pro	Leu	Val	Leu	Ile	Gly	His	Phe	Phe	Ala	Val	Ala	Ile	Tyr	Ala
			515				520						525		
Val	Tyr	Phe	Cys	Phe	Lys	Ser	Glu	Pro	Trp	Ile	Thr	Lys	Pro	Arg	Ala
530						535						540			
Leu	Leu	Ser	Ser	Gly	Ala	Val	Leu	Tyr	Lys	Ala	Cys	Ser	Val	Ile	Phe
545					550					555					560
Pro	Leu	Ile	Tyr	Ser	Glu	Met	Lys	Tyr	Met	Val	His				
				565					570						

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2443 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY:
- (B) CLONE: Consensus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

TTCGCCAGCT CCCCAGGATTG AAGGTTGCCT GGAGCCGCAC TCTTGAGTCC GAGGCCATCT      60
TTTGTGGGAG AAGGCGTCGG CGTTGGCGTT TTCCCCAGGT TGGGCTGTAC AGTGTCTCCG      120
TCCGCGGAAA AAGAAGCCTC TGAACCCGCG CCGGTCCGCA GCCCCCGTGC CTTCCGGACG      180
CTGCTCGCCG TCGCCAGAGG CTAGGCCACG TTTCCCCAG TGTCGAGGTG TTTCTGCGAC      240
CCTCCCTCCA CTCCCATTC CTTCTGAAAG GGCACCTGCT CTTGGTGAGA AAAGAAATTA      300
TCGCACGAAG AGCCAGTATC AGAAGAGTAT CCATCACCCG CAGCAACCGC TCAGGGAACA      360
CCATCAAAAA AGAAAAAAG GGAATATCTG GATTTCCTGG GCGAGGAGGA GCGAGTCTGC      420
TCGGGAGCTG TTCCAGCAGG CGATTTTTAA ATACTGCTTT CTACGCCCTA TACAACCTGG      480
CTTCACATAC TTTTACACTA ACTTTATATG ATTTTAAAA ACTGGTCTGA TCGGACTTCT      540
CGTCCTGGGA CACTGTTTAC TGGAGTCTGG CCGGCTCTCC GTGCTCCTCT TGGTACCTCA      600
TTTGGGGGAG AACCTTAAAC CCACTCGAGC AGATAATCTC CGCCTTGACC GGTGCCACCA      660
AAGAAGCCTT GGAACCATGT GGAATTTTCT GGCATTGCC ACTTTCACCT ATTTTATATA      720
GAAGTTCGGG GACTTCATCA CTTTGGCCAA CAGGGAGGTC CTGTTGTGCG TGCTGGTGTT      780
CCTCTCGCTG GGCCTGGTGC TCTCCTACCG CTGTCGCCAC CGAAACGGGG GTCTCCTCGG      840
GCGCCAGAAG AGCGGCTCCC AGATCGCCCT CTTCTCGGAT ATTCTCTCAG GCCTGCCTTT      900
CATTGGCTTC TTCTGGGCAA ATCCCCCCT GAATCAGAAA ATAAGGAGCA GCTCGAGGCA      960
GGAGGCGCAG AAAAGGAACC AATATTTTCT AAACAAGCTT AATAGGAACA GCTGCTGTAC      1020
ATCAACATCT TCTCAGAATG ACCCAGAAGT TATCATCGTG GGAGCTGGCG TGCTTGGCTC      1080
TGCTTTGGCA GCTGTGCTTT CCAGAGATGG AAGAAAGGTG ACAGTCATTG AGAGAGACTT      1140
AAAAGAGCCT GACAGAATAG TTGGAGAATT CCTGCAGCCG GGTGGTTATC ATGTTCTCAA      1200
AGACCTTGGT CTTGGAGATA CAGTGGAAGG TCTTGATGCC CAGGTTGTAA ATGTTACAT      1260
GATTCATGAT CAGGAAAGCA AATCAGAGGT TCAGATTCCT AACCTCTGT CAGAAAACAA      1320
TCAAGTGCAG AGTGGAAGAG CTTTCCACCA CGGAAGATTC ATCATGAGTC TCCGGAAGGC      1380
AGTTATGGCA GAGCCCAATG CAAAGTTTAT TGAAGGTGTT GTGTTACAGT TATTAGAGGA      1440
AGATGATGTT GTGATGGGAG TTCAGCACAA GGATAAAGAG ACTGGAGATA TCAAGGAAC      1500
CCATGCTCCA CTGACTGTTG TTGCAGATGG GCTTTTCTCC AAGTTCAGGA AAAGCCTGGT      1560
CTCCAATAAA GTTTCTGTAT CATCTCATTT TGTTGGCTTT CTTATGAAGA ATGCACCACA      1620
GTTTACAGCA AATCATGCTG AACTTATTTT AGCTAACCCG AGTCCAGTTC TCATCTACCA      1680
GATTTTATCC AGTGAAATCG AGTACTTGTT GACATTAGAG GGAATGCCAA GGAATTTAAG      1740
AGAATACATG GTTGAAAAAA TTTACCCACA AATACCTGAT CACCTGAAAG AACCATTCTT      1800
AGAAGCCACT GACAATTCTC ATCTGAGGTC CATGCCAGCA AGCTTCCTTC CTCCTTCATC      1860
AGTGAAGAAA CGAGGTGTTT TTCTTTTGGG AGACGCATAT AATATGAGGC ATCCACTTAC      1920
TGGTGGAGGA ATGACTGTTG CTTTTAAAGA TATAAACTA TGGAGAAAAC TGCTAAAGGG      1980
TATCCCTGAC CTTTATGATG ATGCAGCTAT TTTCGAGGCC AAAAATCAT TTTACTGGGC      2040
AAGAAAAACA TCTCATTCCT TTGTCGTGAA TATCCTTGCT CAGGCTCTTT ATGAATTATT      2100
TTCTGCCACA GATGATTCCC TGCATCAACT AAGAAAAGCC TGTTTTCTTT ATTTCAAAC      2160
TGGTGGCGAA TGTNTTGGCG GTCCTGTTGG GCTGCTTCT GTATTGTCTC CTAACCCTCT      2220
AGTTTTAATT GGACACTTCT TTGCTGTTGC AATCTATGCC GTGTATTTT GCTTTAAGTC      2280
AGAACCTTGG ATTACAAAAC CTCGAGCCCT TCTCAGTAGT GGTGCTGTAT TGTACAAAGC      2340
GTGTTCTGTA ATATTTCTCT TAATTTACTC AGAAATGAAG TATATGGTTC ATTAAGCTTA      2400
AAGGGGAACC ATTTGTGAAT GAATATTTGG AACTTACCAA GTC      2443

```

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 572 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vii) IMMEDIATE SOURCE:

(A) LIBRARY: GenBank
(B) CLONE: 1217593

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```

Met Trp Thr Phe Leu Gly Ile Ala Thr Phe Thr Tyr Phe Tyr Lys Lys
 1          5          10          15
Cys Gly Asp Val Thr Leu Ala Asn Lys Glu Leu Leu Leu Cys Val Leu
 20          25          30
Val Phe Leu Ser Leu Gly Leu Val Leu Ser Tyr Arg Cys Arg His Arg
 35          40          45
His Gly Gly Leu Leu Gly Arg His Gln Ser Gly Ala Gln Phe Ala Ala
 50          55          60
Phe Ser Asp Ile Leu Ser Ala Leu Pro Leu Ile Gly Phe Phe Trp Ala
 65          70          75          80
Lys Ser Pro Glu Ser Glu Lys Lys Glu Gln Leu Glu Ser Lys Lys Cys
 85          90          95
Arg Lys Glu Ile Gly Leu Ser Glu Thr Thr Leu Thr Gly Ala Ala Thr
 100          105          110
Ser Val Ser Thr Ser Phe Val Thr Asp Pro Glu Val Ile Ile Val Gly
 115          120          125
Ser Gly Val Leu Gly Ser Ala Leu Ala Ala Val Leu Ser Arg Asp Gly
 130          135          140
Arg Lys Val Thr Val Ile Glu Arg Asp Leu Lys Glu Pro Asp Arg Ile
 145          150          155          160
Val Gly Glu Leu Leu Gln Pro Gly Gly Tyr Arg Val Leu Gln Glu Leu
 165          170          175
Gly Leu Gly Asp Thr Val Glu Gly Leu Asn Ala His His Ile His Gly
 180          185          190
Tyr Ile Val His Asp Tyr Glu Ser Arg Ser Glu Val Gln Ile Pro Tyr
 195          200          205
Pro Leu Ser Glu Thr Asn Gln Val Gln Ser Gly Ile Ala Phe His His
 210          215          220
Gly Arg Phe Ile Met Ser Leu Arg Lys Ala Ala Met Ala Glu Pro Asn
 225          230          235          240
Val Lys Phe Ile Glu Gly Val Val Leu Gln Leu Leu Glu Glu Asp Asp
 245          250          255
Ala Val Ile Gly Val Gln Tyr Lys Asp Lys Glu Thr Gly Asp Thr Lys
 260          265          270
Glu Leu His Ala Pro Leu Thr Val Val Ala Asp Gly Leu Phe Ser Lys
 275          280          285
Phe Arg Lys Ser Leu Ile Ser Ser Lys Val Ser Val Ser Ser His Phe
 290          295          300
Val Gly Phe Leu Met Lys Asp Ala Pro Gln Phe Lys Pro Asn Phe Ala
 305          310          315          320
Glu Leu Val Leu Val Asn Pro Ser Pro Val Leu Ile Tyr Gln Ile Ser
 325          330          335
Ser Ser Glu Thr Arg Val Leu Val Asp Ile Arg Gly Glu Leu Pro Arg
 340          345          350
Asn Leu Arg Glu Tyr Met Ala Glu Gln Ile Tyr Pro Gln Leu Pro Glu
 355          360          365
His Leu Lys Glu Ser Phe Leu Glu Ala Ser Gln Asn Gly Arg Leu Arg
 370          375          380
Thr Met Pro Ala Ser Phe Leu Pro Pro Ser Ser Val Asn Lys Arg Gly
 385          390          395          400
Val Leu Ile Leu Gly Asp Ala Tyr Asn Leu Arg His Pro Leu Thr Gly
 405          410          415
Gly Gly Met Thr Val Ala Leu Lys Asp Ile Lys Leu Trp Arg Gln Leu
 420          425          430
Leu Lys Asp Ile Pro Asp Leu Tyr Asp Asp Ala Ala Ile Phe Gln Ala
 435          440          445
Lys Lys Ser Phe Phe Trp Ser Arg Lys Arg Thr His Ser Phe Val Val
 450          455          460
Asn Val Leu Ala Gln Ala Leu Tyr Glu Leu Phe Ser Ala Thr Asp Asp
 465          470          475          480
Ser Leu His Gln Leu Arg Lys Ala Cys Phe Leu Tyr Phe Lys Leu Gly
 485          490          495

```

Gly	Glu	Cys	Val	Thr	Gly	Pro	Val	Gly	Leu	Leu	Ser	Ile	Leu	Ser	Pro
			500					505					510		
His	Pro	Leu	Val	Leu	Ile	Arg	His	Phe	Phe	Ser	Val	Ala	Ile	Tyr	Ala
		515				520						525			
Thr	Tyr	Phe	Cys	Phe	Lys	Ser	Glu	Pro	Trp	Ala	Thr	Lys	Pro	Arg	Ala
	530					535					540				
Leu	Phe	Ser	Ser	Gly	Ala	Val	Leu	Tyr	Lys	Ala	Cys	Ser	Ile	Leu	Phe
545					550					555					560
Pro	Leu	Ile	Tyr	Ser	Glu	Met	Lys	Tyr	Leu	Val	His				
				565					570						

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 573 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: GenBank
 (B) CLONE: 1083804

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met	Trp	Thr	Phe	Leu	Gly	Ile	Ala	Thr	Phe	Thr	Tyr	Phe	Tyr	Lys	Lys
1				5					10					15	
Cys	Gly	Asp	Val	Thr	Leu	Ala	Asn	Lys	Glu	Leu	Leu	Leu	Cys	Val	Leu
			20				25					30			
Val	Phe	Leu	Ser	Leu	Gly	Leu	Val	Ser	Tyr	Arg	Cys	Arg	His	Arg	
		35				40					45				
Asn	Gly	Gly	Leu	Leu	Gly	Arg	His	Gln	Ser	Gly	Ser	Gln	Phe	Ala	Ala
	50				55					60					
Phe	Ser	Asp	Ile	Leu	Ser	Ala	Leu	Pro	Leu	Ile	Gly	Phe	Phe	Trp	Ala
65				70					75					80	
Lys	Ser	Pro	Pro	Glu	Ser	Glu	Lys	Lys	Glu	Gln	Leu	Glu	Ser	Lys	Arg
			85					90					95		
Arg	Arg	Lys	Glu	Val	Asn	Leu	Ser	Glu	Thr	Thr	Leu	Thr	Gly	Ala	Ala
			100				105					110			
Thr	Ser	Val	Ser	Thr	Ser	Ser	Val	Thr	Asp	Pro	Glu	Val	Ile	Ile	Ile
		115				120					125				
Gly	Ser	Gly	Val	Leu	Gly	Ser	Ala	Leu	Ala	Thr	Val	Leu	Ser	Arg	Asp
	130				135						140				
Gly	Arg	Thr	Val	Thr	Val	Ile	Glu	Arg	Asp	Leu	Lys	Glu	Pro	Asp	Arg
145				150					155					160	
Ile	Leu	Gly	Glu	Cys	Leu	Gln	Pro	Gly	Gly	Tyr	Arg	Val	Leu	Arg	Glu
			165					170					175		
Leu	Gly	Leu	Gly	Asp	Thr	Val	Glu	Ser	Leu	Asn	Ala	His	His	Ile	His
		180				185					190				
Gly	Tyr	Val	Ile	His	Asp	Cys	Glu	Ser	Arg	Ser	Glu	Val	Gln	Ile	Pro
	195					200					205				
Tyr	Pro	Val	Ser	Glu	Asn	Asn	Gln	Val	Gln	Ser	Gly	Val	Ala	Phe	His
	210				215						220				
His	Gly	Lys	Phe	Ile	Met	Ser	Leu	Arg	Lys	Ala	Ala	Met	Ala	Glu	Pro
225				230					235					240	
Asn	Val	Lys	Phe	Ile	Glu	Gly	Val	Val	Leu	Arg	Leu	Leu	Glu	Glu	Asp
			245					250					255		
Asp	Ala	Val	Ile	Gly	Val	Gln	Tyr	Lys	Asp	Lys	Glu	Thr	Gly	Asp	Thr
		260				265						270			
Lys	Glu	Leu	His	Ala	Pro	Leu	Thr	Val	Val	Ala	Asp	Gly	Leu	Phe	Ser
	275					280					285				
Lys	Phe	Arg	Lys	Asn	Leu	Ile	Ser	Asn	Lys	Val	Ser	Val	Ser	Ser	His
	290				295						300				
Phe	Val	Gly	Phe	Ile	Met	Lys	Asp	Ala	Pro	Gln	Phe	Lys	Ala	Asn	Phe
305				310					315						320

Ala Glu Leu Val Leu Val Asp Pro Ser Pro Val Leu Ile Tyr Gln Ile
 325 330
 Ser Pro Ser Glu Thr Arg Val Leu Val Asp Ile Arg Gly Glu Leu Pro
 340 345 350
 Arg Asn Leu Arg Glu Tyr Met Thr Glu Gln Ile Tyr Pro Gln Ile Pro
 355 360 365
 Asp His Leu Lys Glu Ser Phe Leu Glu Ala Cys Gln Asn Ala Arg Leu
 370 375 380
 Arg Thr Met Pro Ala Ser Phe Leu Pro Pro Ser Ser Val Asn Lys Arg
 385 390 395 400
 Gly Val Leu Leu Leu Gly Asp Ala Tyr Asn Leu Arg His Pro Leu Thr
 405 410 415
 Gly Gly Gly Met Thr Val Ala Leu Lys Asp Ile Lys Ile Trp Arg Gln
 420 425 430
 Leu Leu Lys Asp Ile Pro Asp Leu Tyr Asp Asp Ala Ala Ile Phe Gln
 435 440 445
 Ala Lys Lys Ser Phe Phe Trp Ser Arg Lys Arg Ser His Ser Phe Val
 450 455 460
 Val Asn Val Leu Ala Gln Ala Leu Tyr Glu Leu Phe Ser Ala Thr Asp
 465 470 475 480
 Asp Ser Leu Arg Gln Leu Arg Lys Ala Cys Phe Leu Tyr Phe Lys Leu
 485 490 495
 Gly Gly Glu Cys Leu Thr Gly Pro Val Gly Leu Leu Ser Ile Leu Ser
 500 505 510
 Pro Asp Pro Leu Leu Leu Ile Arg His Phe Phe Ser Val Ala Val Tyr
 515 520 525
 Ala Thr Tyr Phe Cys Phe Lys Ser Glu Pro Trp Ala Thr Lys Pro Arg
 530 535 540
 Ala Leu Phe Ser Ser Gly Ala Ile Leu Tyr Lys Ala Cys Ser Ile Ile
 545 550 555 560
 Phe Pro Leu Ile Tyr Ser Glu Met Lys Tyr Leu Val His
 565 570

What is claimed is:

1. A substantially purified squalene epoxidase comprising the amino acid sequence of SEQ ID NO:1 or fragments thereof.
2. An isolated and purified polynucleotide sequence encoding the squalene epoxidase of claim 1.
3. A polynucleotide sequence which hybridizes under stringent conditions to the polynucleotide sequence of claim 2.
4. A hybridization probe comprising the polynucleotide sequence of claim 2.
5. An isolated and purified polynucleotide sequence comprising SEQ ID NO:2 or variants thereof.
6. A polynucleotide sequence which is complementary to SEQ ID NO:2 or variants thereof.
7. A hybridization probe comprising the polynucleotide sequence of claim 6.
8. An expression vector containing the polynucleotide sequence of claim 2.
9. A host cell containing the vector of claim 8.
10. A method for producing a polypeptide comprising a polypeptide of SEQ ID NO:1 the method comprising the steps of:
 - a) culturing the host cell of claim 9 under conditions suitable for the expression of the polypeptide; and
 - b) recovering the polypeptide from the host cell culture.
11. A purified antibody which binds specifically to the polypeptide of claim 1.
12. A purified agonist which specifically binds to and modulates the activity of the polypeptide of claim 1.
13. A purified antagonist which specifically binds to and blocks the activity of the polypeptide of claim 1.
14. A pharmaceutical composition comprising the antagonist of squalene epoxidase of claim 13 in conjunction with a suitable pharmaceutical carrier.
15. A method for treating hypercholesterolemia comprising administering to a subject in need of such treatment an effective amount of the composition of claim 14.

16. A method for detection of polynucleotides encoding squalene epoxidase of claim 1 in a biological sample comprising the steps of:

- a) hybridizing a polynucleotide consisting of SEQ ID NO:2 to nucleic acid material of a biological sample, thereby forming a hybridization complex; and
- 5 b) detecting said hybridization complex, wherein the presence of said complex correlates with the presence of a polynucleotide encoding squalene epoxidase in said biological sample.

17. The method of claim 16 wherein before hybridization, the nucleic acid material of the biological sample is amplified by the polymerase chain reaction.

1/13

5'	9	18	27	36	45	54
T	TCG CCA GCT CCC CGG ATT GAA GGT TGC CTG GAG CCG CAC TCT TGA GTC CGA					
63	72	81	90	99	108	
GGC	CAT CTT TTG TTG GAG AAG GCG TCG GCG TTG GCG TTT TCC CGA GGT TGG GCT					
117	126	135	144	153	162	
GTA	CAG TGT CTC CGT CCG CGG AAA AAG AAG CCT CTG AAC CCG CGG TCC GCA					
171	180	189	198	207	216	
GCC	CCC GTG CCT TCC GGA CGC TGC TCG CCG TCG CCA GAG GCT AGG CCA CGT TTC					
225	234	243	252	261	270	
CCC	CAG TGT CGA GGT GTT TCT GCG ACC CTC CCT CCA CTC CCA TTC CCT TCT GAA					
279	288	297	306	315	324	
AGG	GCA CCT GCT CTT GGT GAG AAA AGA AAT TAT CGC ACG AAG AGC CAG TAT CAG					
333	342	351	360	369	378	
AAG	AGT ATC CAT CAC CCG CAG CAA CCG CTC AGG GAA CAC CAT CAA AAA AGA AAA					
387	396	405	414	423	432	
AAA	GGG AAT ATC TGG ATT TCC TGG GCG AGG AGG AGC GAG TCT GCT CGG GAG CTG					
441	450	459	468	477	486	
TTC	CAG CAG GCG ATT TTT AAA TAC TGC TTT CTA CGC CCT ATA CAA CTT GGC TTC					

FIGURE 1A

2/13

495	504	513	522	531	540
ACA TAC TTT TAC ACT AAC TTT ATA TGA TTT TTA AAA ACT GGT CTG ATC GGA CTT					
549	558	567	576	585	594
CTC GTC CTG GGA CAC TGT TTA CTG GAG TCT GGC CGG CTC TCC GTG CTC CTC TTG					
603	612	621	630	639	648
GTA CCT CAT TTG GGG GAG AAC CTT AAA CCC ACT CGA GCA GAT AAT CTC CGC CTT					
657	666	675	684	693	702
GAC CGG TGC CAC CAA AGA AGC CTT GGA ACC ATG TGG ACT TTT CTG GGC ATT GCC					
			M W T F L G I A		
711	720	729	738	747	756
ACT TTC ACC TAT TTT TAT AAG AAG TTC GGG GAC TTC ATC ACT TTG GCC AAC AGG					
T F T Y F Y K K F G G D F I T L A N R					
765	774	783	792	801	810
GAG GTC CTG TTG TGC GTG CTG GTG TTC CTC TCG CTG GGC CTG GTG CTC TCC TAC					
E V L L C V L L V F F L S L G L V L S Y					
819	828	837	846	855	864
CGC TGT CGC CAC CGA AAC GGG GGT CTC CTC GGG CGC CAG AAG AGC GGC TCC CAG					
R C R H R N G G L L G R Q K S G S Q					

FIGURE 1B

3/13

873	882	891	900	909	918
ATC GCC CTC TTC TCG GAT ATT CTC TCA GGC CTG CCT TTC ATT GGC TTC TTC TGG					
I A L F S D I L S G L P F I G F W					
927	936	945	954	963	972
GCA AAT CCC CCC CTG AAT CAG AAA ATA AGG AGC AGC TCG AGG CAG GAG GCG CAG					
A N P P L N Q K I R S S S R Q E A Q					
981	990	999	1008	1017	1026
AAA AGG AAC CAA TAT TTC AGA AAC AAG CTT AAT AGG AAC AGC TGC TGT ACA TCA					
K R N Q Y F R N K L N R N S C C T S					
1035	1044	1053	1062	1071	1080
ACA TCT TCT CAG AAT GAC CCA GAA GTT ATC ATC GTG GGA GCT GGC GTG CTT GGC					
T S S Q N D P E V I I V G A G V L G					
1089	1098	1107	1116	1125	1134
TCT GCT TTG GCA GCT GTG CTT TCC AGA GAT GGA AGA AAG GTG ACA GTC ATT GAG					
S A L A A V L S R D G R K V T V I E					
1143	1152	1161	1170	1179	1188
AGA GAC TTA AAA GAG CCT GAC AGA ATA GTT GGA GAA TTC CTG CAG CCG GGT GGT					
R D L K E P D R I V G E F L Q P G G					
1197	1206	1215	1224	1233	1242
TAT CAT GTT CTC AAA GAC CTT GGT CTT GGA GAT ACA GTG GAA GGT CTT GAT GCC					
Y H V L L K D L G L G D T V E G L D A					

FIGURE 1C

1251 1260 1269 1278 1287 1296
 CAG GTT GTA AAT GGT TAC ATG ATT CAT GAT CAG GAA AGC AAA TCA GAG GTT CAG
 Q V V N G Y M I H D Q E S K S E V Q

1305 1314 1323 1332 1341 1350
 ATT CCT AAC CCT CTG TCA GAA AAC AAT CAA GTG CAG AGT GGA AGA GCT TTC CAC
 I P N P L S E N N Q V Q S G R A F H

1359 1368 1377 1386 1395 1404
 CAC GGA AGA TTC ATC ATG AGT CTC CGG AAA GCA GTT ATG GCA GAG CCC AAT GCA
 H G R F I M S L R K A V M A E P N A

1413 1422 1431 1440 1449 1458
 AAG TTT ATT GAA GGT GTT GTG TTA CAG TTA TTA GAG GAA GAT GAT GTT GTG ATG
 K F I E G V V L Q L L E E D D V V M

1467 1476 1485 1494 1503 1512
 GGA GTT CAG CAC AAG GAT AAA GAG ACT GGA GAT ATC AAG GAA CTC CAT GCT CCA
 G V Q H K D K E T G D I K E L H A P

1521 1530 1539 1548 1557 1566
 CTG ACT GTT GTT GCA GAT GGG CTT TTC TCC AAG TTC AGG AAA AGC CTG GTC TCC
 L T V V A D G G L F S K F R K S L V S

1575 1584 1593 1602 1611 1620
 AAT AAA GTT TCT GTA TCA TCT CAT TTT GTT GGC TTT CTT ATG AAG AAT GCA CCA
 N K V S S S H F V G F L M K N A P

FIGURE 1D

5/13

1629	1638	1647	1656	1665	1674
CAG TTT ACA GCA AAT CAT GCT GAA CTT ATT TTA GCT AAC CCG AGT CCA GTT CTC					
Q F T A A N H A E L I L A N P S P V L					
1683	1692	1701	1710	1719	1728
ATC TAC CAG ATT TCA TCC AGT GAA ATC GAG TAC TTG TTG ACA TTA GAG GGA ATG					
I Y Q I S S S E I E Y L L T L E G M					
1737	1746	1755	1764	1773	1782
CCA AGG AAT TTA AGA GAA TAC ATG GTT GAA AAA ATT TAC CCA CAA ATA CCT GAT					
P R N L R E Y M V E K I Y P Q I P D					
1791	1800	1809	1818	1827	1836
CAC CTG AAA GAA CCA TTC TTA GAA GCC ACT GAC AAT TCT CAT CTG AGG TCC ATG					
H L K E P F L L E A T D N S H L R S M					
1845	1854	1863	1872	1881	1890
CCA GCA AGC TTC CTT CCT CCT TCA TCA GTG AAG AAA CGA GGT GTT CTT CTT TTG					
P A S F L P P P S S V K K R G V L L L					
1899	1908	1917	1926	1935	1944
GGA GAC GCA TAT AAT ATG AGG CAT CCA CTT ACT GGT GGA GGA ATG ACT GTT GCT					
G D A Y N M R H P L T G G G M T V A					
1953	1962	1971	1980	1989	1998
TTT AAA GAT ATA AAA CTA TGG AGA AAA CTG CTA AAG GGT ATC CCT GAC CTT TAT					
F K D I K L W R K L L K G I P D L Y					

FIGURE 1E

6/13

2007	2016	2025	2034	2043	2052
GAT GAT GCA GCT ATT TTC GAG GCC AAA AAA TCA TTT TAC TGG GCA AGA AAA ACA					
D D A A I F E A K K S F Y W A R K T					
2061	2070	2079	2088	2097	2106
TCT CAT TCC TTT GTC GTG AAT ATC CTT GCT CAG GCT CTT TAT GAA TTA TTT TCT					
S H S F V V N I L A Q A L Y E L F S					
2115	2124	2133	2142	2151	2160
GCC ACA GAT GAT TCC CTG CAT CAA CTA AGA AAA GCC TGT TTT CTT TAT TTC AAA					
A T D D S L H Q L R K A C F L Y F K					
2169	2178	2187	2196	2205	2214
CTT GGT GGC GAA TGT NTT GCG GGT CCT GTT GGG CTG CTT TCT GTA TTG TCT CCT					
L G G E C X A G P V G L L S V L S P					
2223	2232	2241	2250	2259	2268
AAC CCT CTA GTT TTA ATT GGA CAC TTC TTT GCT GTT GCA ATC TAT GCC GTG TAT					
N P L V L I G H F F A V A I Y A V Y					
2277	2286	2295	2304	2313	2322
TTT TGC TTT AAG TCA GAA CCT TGG ATT ACA AAA CCT CGA GCC CTT CTC AGT AGT					
F C F K S E P W I T K P R A L L S S					
2331	2340	2349	2358	2367	2376
GGT GCT GTA TTG TAC AAA GCG TGT TCT GTA ATA TTT CCT CTA ATT TAC TCA GAA					
G A V L Y K A C S V I F P L I Y S E					

FIGURE 1F

2385 2394 2403 2412 2421 2430
ATG AAG TAT ATG GTT CAT TAA GCT TAA AGG GGA ACC ATT TGT GAA TGA ATA TTT
M K Y M V H

2439
GGA ACT TAC CAA GTC 3'

FIGURE 1G

1	M	W	T	F	L	G	I	A	T	F	T	Y	F	Y	K	K	F	G	D	F	I	T	L	A	N	R	E	V	L	L	C	V	L	V	F	L	S	L	G	L	HSQEP
1	M	W	T	F	L	G	I	A	T	F	T	Y	F	Y	K	K	C	G	D	-	V	T	L	A	N	K	E	L	L	L	C	V	L	V	F	L	S	L	G	L	GI 1217593
1	M	W	T	F	L	G	I	A	T	F	T	Y	F	Y	K	K	C	G	D	-	V	T	L	A	N	K	E	L	L	L	C	V	L	V	F	L	S	L	G	L	GI 1083804
41	V	L	S	Y	R	C	R	H	R	N	G	G	L	L	G	R	Q	K	S	G	S	Q	I	A	L	F	S	D	I	L	S	G	L	P	F	I	G	F	F	W	HSQEP
40	V	L	S	Y	R	C	R	H	R	H	G	G	L	L	G	R	H	Q	S	G	A	Q	F	A	A	F	S	D	I	L	S	A	L	P	L	I	G	F	F	W	GI 1217593
40	V	L	S	Y	R	C	R	H	R	N	G	G	L	L	G	R	H	Q	S	G	S	Q	F	A	A	F	S	D	I	L	S	A	L	P	L	I	G	F	F	W	GI 1083804
81	A	N	-	P	P	L	N	Q	K	I	R	S	S	S	R	Q	E	A	Q	K	R	N	Q	Y	F	R	N	K	L	N	R	N	S	C	C	T	S	T	S	S	HSQEP
80	A	K	S	P	-	E	S	E	K	K	E	Q	L	E	S	K	K	C	R	K	E	I	G	L	S	E	T	T	L	T	G	A	A	T	S	V	S	T	S	F	GI 1217593
80	A	K	S	P	P	E	S	E	K	K	E	Q	L	E	S	K	R	R	K	E	V	N	L	S	E	T	T	L	T	G	A	A	T	S	V	S	T	S	S	GI 1083804	
120	Q	N	D	P	E	V	I	I	V	G	A	G	V	L	G	S	A	L	A	A	V	L	S	R	D	G	R	K	V	T	V	I	E	R	D	L	K	E	P	D	HSQEP
119	V	T	D	P	E	V	I	I	V	G	S	G	V	L	G	S	A	L	A	A	V	L	S	R	D	G	R	K	V	T	V	I	E	R	D	L	K	E	P	D	GI 1217593
120	V	T	D	P	E	V	I	I	T	G	S	G	V	L	G	S	A	L	A	T	V	L	S	R	D	G	R	T	V	T	V	I	E	R	D	L	K	E	P	D	GI 1083804
160	R	I	V	G	E	F	L	Q	P	G	G	Y	H	V	L	K	D	L	G	L	G	D	T	V	E	G	L	D	A	Q	V	V	N	G	Y	M	I	H	D	Q	HSQEP
159	R	I	V	G	E	L	L	Q	P	G	G	Y	R	V	L	Q	E	L	G	L	G	D	T	V	E	G	L	N	A	H	H	I	H	G	Y	I	V	H	D	Y	GI 1217593
160	R	I	L	G	E	C	L	Q	P	G	G	Y	R	V	L	R	E	L	G	L	G	D	T	V	E	S	L	N	A	H	H	I	H	G	Y	V	I	H	D	C	GI 1083804
200	E	S	K	S	E	V	Q	I	P	N	P	L	S	E	N	N	Q	V	Q	S	G	R	A	F	H	H	G	R	F	I	M	S	L	R	K	A	V	M	A	E	HSQEP
199	E	S	R	S	E	V	Q	I	P	Y	P	L	S	E	N	N	Q	V	Q	S	G	I	A	F	H	H	G	R	F	I	M	S	L	R	K	A	A	M	A	E	GI 1217593
200	E	S	R	S	E	V	Q	I	P	Y	P	V	S	E	N	N	Q	V	Q	S	G	V	A	F	H	H	G	K	F	I	M	S	L	R	K	A	A	M	A	E	GI 1083804

FIGURE 2A

240	P N A K F I E G V V L Q L L E E D D V V M G V Q Q H K D K E T G D I K E L H A P L	HSQEP
239	P N V K F I E G V V L Q L L E E D D A V I G V Q Q Y K D K E T G D T K E L H A P L	GI 1217593
240	P N V K F I E G V V L R L L E E D D A V I G V Q Q Y K D K E T G D T K E L H A P L	GI 1083804
280	T V V A D G L F S K F R K S L V S N K V S V S S H F V G F L M K N A P Q F T A N	HSQEP
279	T V V A D G L F S K F R K S L I S K V S V S S H F V G F L M K D A P Q F K P N	GI 1217593
280	T V V A D G L F S K F R K N L I S N K V S V S S H F V G F I M K D A P Q F K A N	GI 1083804
320	H A E L I L A N P S P V L I Y Q I S S S E I E Y L L T L E G - M P R N L R E Y M	HSQEP
319	F A E L V L V N P S P V L I Y Q I S S S E T R V L V D I R G E L P R N L R E Y M	GI 1217593
320	F A E L V L V D P S P V L I Y Q I S P S E T R V L V D I R G E L P R N L R E Y M	GI 1083804
359	V E K I Y P Q I P D H L K E P F L E A T D N S H L R S M P A S F L P P S S V K K	HSQEP
359	A E Q I Y P Q I P E H L K E S F L E A S Q N G R L R T M P A S F L P P S S V N K	GI 1217593
360	T E Q I Y P Q I P D H L K E S F L E A C Q N A R L R T M P A S F L P P S S V N K	GI 1083804
399	R G V L L L G D A Y N M R H P L T G G G M T V A F K D I K L W R K L L K G I P D	HSQEP
399	R G V L L L G D A Y N L R H P L T G G G M T V A L K D I K L W R Q L L K D I P D	GI 1217593
400	R G V L L L G D A Y N L R H P L T G G G M T V A L K D I K I W R Q L L K D I P D	GI 1083804
439	L Y D D A A I F E A K K S F Y W A R K T S H S F V V N I L A Q A L Y E L F S A T	HSQEP
439	L Y D D A A I F Q A K K S F F W S R K R T H S F V V N V L A Q A L Y E L F S A T	GI 1217593
440	L Y D D A A I F Q A K K S F F W S R K R S H S F V V N V L A Q A L Y E L F S A T	GI 1083804

FIGURE 2B

479	D	D	S	L	H	Q	L	R	K	A	C	F	L	Y	F	K	L	G	G	E	C	X	A	G	P	V	G	L	L	S	V	L	S	P	N	P	L	V	L	I	HSQEP
479	D	D	S	L	H	Q	L	R	K	A	C	F	L	Y	F	K	L	G	G	E	C	V	T	G	P	V	G	L	L	S	I	L	S	P	H	P	L	V	L	I	GI 1217593
480	D	D	S	L	R	Q	L	R	K	A	C	F	L	Y	F	K	L	G	G	E	C	L	T	G	P	V	G	L	L	S	I	L	S	P	D	P	L	L	I	GI 1083804	
519	G	H	F	F	A	V	A	I	Y	A	V	Y	F	C	F	K	S	E	P	W	I	T	K	P	R	A	L	L	S	S	G	A	V	L	Y	K	A	C	S	V	HSQEP
519	R	H	F	F	S	V	A	I	Y	A	T	Y	F	C	F	K	S	E	P	W	A	T	K	P	R	A	L	F	S	S	G	A	V	L	Y	K	A	C	S	I	GI 1217593
520	R	H	F	F	S	V	A	V	Y	A	T	Y	F	C	F	K	S	E	P	W	A	T	K	P	R	A	L	F	S	S	G	A	I	L	Y	K	A	C	S	I	GI 1083804
559	I	F	P	L	I	Y	S	E	M	K	Y	M	V	H																											
559	L	F	P	L	I	Y	S	E	M	K	Y	L	V	H																											
560	I	F	P	L	I	Y	S	E	M	K	Y	L	V	H																											

FIGURE 2C




FIGURE 3

12/13




FIGURE 4




FIGURE 5

INTERNATIONAL SEARCH REPORT

Intern: al Application No

PCT/US 97/20681

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/53 C12N9/02 C12Q1/68 C12N1/21 C07K16/40
A61K38/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	NAKAMURA, YUICHI ET AL: "Transcriptional regulation of squalene epoxidase by sterols and inhibitors in HeLa cells" J. BIOL. CHEM. (1996), 271(14), 8053-6 CODEN: JBCHA3; ISSN: 0021-9258, XP002058735 see the whole document ---	1
Y	DATABASE WPI Section Ch, Week 9539 Derwent Publications Ltd., London, GB; Class B04, AN 95-298148 XP002058737 & JP 07 194 381 A (BANYU PHARM CO LTD) , 1 August 1995 see abstract --- -/--	1

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

13 March 1998

Date of mailing of the international search report

27.03.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Delanghe, L

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/20681

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>JUN SAKAKIBARA ET AL.: "Molecular cloning and expression of rat squalene epoxidase" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 1, 6 January 1995, MD US, pages 17-20, XP002058736 see the whole document -----</p>	1

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 97/20681

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 97/20681

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

Claims Nos.: 15

because they relate to subject matter not required to be searched by this Authority, namely:

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

Remark : Although claim 15 is directed to a method of treatment of the human/animal body , the search has been carried out and based on the alleged effects of the compound/composition.





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/53, 9/02, C12Q 1/68, C12N 1/21, C07K 16/40, A61K 38/44	A1	(11) International Publication Number: WO 98/20137 (43) International Publication Date: 14 May 1998 (14.05.98)
(21) International Application Number: PCT/US97/20681 (22) International Filing Date: 5 November 1997 (05.11.97) (30) Priority Data: 08/745,934 7 November 1996 (07.11.96) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 08/745,934 (CIP) Filed on 7 November 1996 (07.11.96) (71) Applicant (for all designated States except US): INCTYE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): HILLMAN, Jennifer, L. [US/US]; 230 Monroe Drive #12, Mountain View, CA 94040 (US). HAWKINS, Phillips, R. [US/US]; 750 N. Shoreline Boulevard #96, Mountain View, CA 94034 (US). (74) Agent: BILLINGS, Lucy, J.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US).	(81) Designated States: AT, AU, BR, CA, CH, CN, DE, DK, ES, FI, GB, IL, JP, KR, MX, NO, NZ, RU, SE, SG, US, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: HUMAN SQUALENE EPOXIDASE		
(57) Abstract <p>The present invention provides a human squalene epoxidase (HSQEP) and polynucleotides which identify and encode HSQEP. The invention also provides genetically engineered expression vectors and host cells comprising the nucleic acid sequences encoding HSQEP and a method for producing HSQEP. The invention also provides for the use of HSQEP and agonists, antibodies, or antagonists specifically binding HSQEP, in the prevention and treatment of diseases associated with expression of HSQEP. Additionally, the invention provides for the use of antisense molecules to polynucleotides encoding HSQEP for the treatment of diseases associated with the expression of HSQEP. The invention also provides diagnostic assays which utilize the polynucleotide, or fragments or the complement thereof, and antibodies specifically binding HSQEP.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		